

**Women who quit smoking during pregnancy and relapse in  
the first two years after pregnancy.**

A Thesis

by

Priya Srikanth

Presented to the Department of Public Health and Preventive Medicine  
Oregon Health and Science University  
in partial fulfillment of  
the requirements for the degree of  
Master of Public Health  
May 2009



Department of Public Health and Preventive Medicine

School of Medicine

Oregon Health & Science University

---

CERTIFICATE OF APPROVAL

---

This is to certify that the Master's thesis of

Priya Srikanth

has been approved

---

Kenneth D. Rosenberg, MD, MPH

---

*Hedi Lapidus, PhD*

---

*Elizabeth Adams, PhD, RD*

# TABLE OF CONTENTS

LIST OF TABLES.....	ii
LIST OF ABBREVIATIONS.....	iii
ACKNOWLEDGEMENTS.....	iv
ABSTRACT.....	v
INTRODUCTION	
QUITTING SMOKING DURING PREGNANCY.....	1
STAYING QUIT, 2 YEARS POSTPARTUM.....	3
SPECIFIC AIMS.....	4
METHODS	
PRAMS.....	6
PRAMS-2.....	7
VARIABLE CODING.....	10
STATISTICAL ANALYSIS.....	26
RESULTS	
SUMMARY.....	28
QUIT SMOKING DURING PREGNANCY: UNIVARIATE ANALYSIS.....	31
QUIT SMOKING DURING PREGNANCY: MULTIVARIATE ANALYSIS.....	37
STAYED QUIT, 2 YEARS POSTPARTUM: UNIVARIATE ANALYSIS.....	40
STAYED QUIT, 2 YEARS POSTPARTUM: MULTIVARIATE ANALYSIS.....	46
DISCUSSION	
SUMMARY.....	50
QUITTING SMOKING DURING PREGNANCY.....	51
STAYING QUIT, 2 YEARS POSTPARTUM.....	53
STRENGTHS AND LIMITATIONS.....	54
PUBLIC HEALTH IMPLICATIONS.....	58
FUTURE STUDIES.....	60
SUMMARY AND CONCLUSION.....	61
APPENDIX A.....	62
APPENDIX B.....	63
APPENDIX C.....	64
REFERENCES.....	65

## **List of Tables**

Table 1	Coding of quitting smoking during pregnancy outcome variable
Table 2	Patterns of smoking among women who reported smoking at least 100 cigarettes in the past 2 years
Table 3	Independent variables derived from Birth certificate
Table 4	Stressful Life Event Measures, 2004 Oregon PRAMS
Table 5	Independent variables derived from 2004 Oregon PRAMS (for the outcome of quitting smoking during pregnancy)
Table 6	Independent variables derived from 2004 Oregon PRAMS (for the outcome of staying quit 2 years postpartum)
Table 7	Social Support Measures from 2006 Oregon PRAMS-2
Table 8	Independent variables derived from 2006 Oregon PRAMS-2 (used for the outcome of staying quit 2 years postpartum)
Table 9	Risk factors for quitting smoking during pregnancy
Table 10	Multivariable model – risk factors for quitting smoking during pregnancy
Table 11	Forward manual stepwise – multivariable logistic regression model, odds ratios and p-values for quitting smoking during pregnancy
Table 12	Risk factors for staying quit, 2 years postpartum
Table 13	Multivariable model – risk factors for staying quit, 2 years postpartum
Table 14	Forward manual stepwise - multivariable logistic regression model, odds ratios and p-values for staying quit, 2 years postpartum
Table 15	Characteristics of PRAMS-2 respondents versus those lost to follow-up

## **List of Abbreviations**

FPL	Federal Poverty Level
IPV	Intimate Partner Violence
HCW	Health Care Worker
OR	Odds Ratio
WIC	Special Supplementary Nutrition Program for Women, Infants, and Children
PRAMS	Pregnancy Risk Assessment Monitoring System
CI	Confidence Interval
CDC	Centers for Disease Control and Prevention
SIDS	Sudden Infant Death Syndrome
NICU	Neonatal Intensive Care Unit
SLE	Stressful Life Event
SHS	Second-hand Smoke

## **Acknowledgements**

First, I would like to thank the members of my thesis committee for their time and support on this endeavor. My thesis chair, Ken Rosenberg, was instrumental in guiding me through my internship, helping me select a thesis topic and in providing me with the Oregon PRAMS/PRAMS-2 dataset. He has always taken time out to give me precise and constructive feedback on all issues or concerns that I had during my internship and my thesis work. Jodi Lapidus guided me through my analysis and helped me understand and interpret output that sometimes was quite intimidating apart from ensuring that I stayed focused and on schedule. Elizabeth Adams was supportive through my entire thesis project and provided me with insightful feedback as needed.

Next, I would like to thank my husband, Srikanth T. Srinivasan, without whose amazing support and patience, fulfilling my goal of obtaining the M.P.H. degree would have been impossible. I would also like to thank my daughter – Anagha Srikanth – and son – Krishna Srikanth – who, in their own way, were my champions and supporters.

I extend my gratitude to Al Sandoval at Oregon Public Health Division for making the PRAMS/PRAMS-2 dataset available to me and for clarifying all data issues as and when I brought them up and to Tina Kent for information on PRAMS and PRAMS-2 methodology.

Appreciation must go to Mike Lasarev for providing me with some technical pointers and for always willing to answer my thesis-related questions.

Finally, many thanks to the faculty of Oregon Health and Science University, department of Public Health and Preventive Medicine, for giving me this opportunity to further enhance my education and knowledge.

## **Abstract**

### Background

The health consequences of smoking during pregnancy, both to the prospective mother and to the infant, have long been recognized. However, very little is known about why some women avoid detrimental behaviors and engage in beneficial ones, while others do not. It is encouraging to note that approximately 40-44% of women who smoke at the start of pregnancy quit smoking for the duration of the pregnancy. Unfortunately, more than half of these women return to smoking within 6 months postpartum. While trends in smoking prevalence during pregnancy are known, little is known about trends in quitting during pregnancy and staying quit, postpartum.

This project aims to identify and differentiate between characteristics of those women who quit smoking during pregnancy and managed to stay quit, 2 years postpartum. Women who smoked during pregnancy and those who quit smoking during pregnancy will be compared to identify characteristics of women who quit smoking during pregnancy. In addition, we aim to identify characteristics of those women who stayed quit, 2 years postpartum compared to those women who relapsed into smoking, 2 years postpartum.

### Methods

This study used data from the 2004 Oregon Pregnancy Risk Assessment Monitoring System (PRAMS) and its 2006 longitudinal follow-up, PRAMS-2. 1,968 women were surveyed with the 2004 Oregon PRAMS and 865 of these women responded to the 2-year follow-up survey in 2006 with a weighted response rate of 51.1%.



In the PRAMS questionnaire, women were asked: “Have you smoked 100 cigarettes in the past 2 years? (A pack has 20 cigarettes.)” Only those who reported smoking at least 100 cigarettes in the past 2 years were included in this analysis. Smoking intensity was measured at three time points in PRAMS – 3 months before pregnancy (T1), last 3 months of pregnancy (T2) and 2-6 months postpartum (T3) with the question: “In the (time point), how many cigarettes did you smoke on an average day? (A pack has 20 cigarettes.)” PRAMS-2 measured smoking intensity at a fourth time point – 2 years postpartum (T4) – only to those who answered ‘Yes’ to the question: “Have you smoked 100 cigarettes in your entire life? (A pack has 20 cigarettes.)” Possible responses were *None, Less than 1 cigarette, 1 to 5 cigarettes, 6 to 10 cigarettes, 11 to 20 cigarettes, 21 to 40 cigarettes, 41 cigarettes or more.*

The respondents were classified into the following categories: (i) Smoked during pregnancy, (ii) Quit smoking during pregnancy, (iii) Stayed quit 2 years postpartum, (iv) Relapsed into smoking, 2 years postpartum, and, (v) Persistent smokers – Women who smoked at all the 4 time points.

Women who smoked during pregnancy were compared with women who quit smoking during pregnancy to identify characteristics of those women who quit smoking during pregnancy. Women who stayed quit 2 years postpartum were compared with women who relapsed into smoking 2 years postpartum to explore characteristics of women who stayed quit 2 years postpartum.

Logistic regression was used to highlight characteristics of women who quit smoking during pregnancy and identify potential correlates of staying quit, 2 years postpartum. Hosmer & Lemeshow’s model building techniques were used to build a

multivariable logistic regression model to determine the characteristics of women who quit smoking during pregnancy and of women who stayed quit, 2 years postpartum. All analyses used weighted data to account for the complex sampling design utilized by PRAMS and PRAMS-2.

### **Results - Quit smoking during pregnancy**

In logistic regression analyses, marginally significant factors associated with quitting smoking during pregnancy were annual household income during pregnancy and WIC participation during pregnancy. Even though these results were only marginally significant in univariate analysis, the magnitude of the Odds Ratios warrants discussion of these results for public health implications.

**Annual household income during pregnancy:** Women who had higher annual household incomes ( $\geq 185\%$  FPL category) were more likely to quit smoking than those who had lower annual household incomes (0-184% FPL category) (Unadjusted OR 3.68; 95% CI 0.98, 13.79), (Adjusted OR 2.17; 95% CI 0.50, 9.42).

**WIC during pregnancy:** Women who did not use WIC during pregnancy were more likely to quit smoking during pregnancy than women who used WIC (Unadjusted OR 2.84; 95% CI 0.87, 9.21), (Adjusted OR 1.08; 95% CI 0.29, 4.03).

### **Results - Stayed quit, 2 years postpartum**

Significant factors associated with women staying quit, 2 years postpartum, were:

**Maternal Education:** Women who had greater than a high school education were more likely to stay quit, 2 years postpartum, than women who had a high school education or less (Unadjusted OR 20.33; 95% CI 4, 103.52), (Adjusted OR 13.30; 95% CI 2.36, 74.87).

**Depressive Symptoms in 13-24 months after delivery:** This was perhaps the most striking finding of this study. Women who reported depressive symptoms 13-24 months after delivery were more likely to stay quit than women who did not report any depressive symptoms in the past 12 months (Unadjusted OR: 6.73, 95% CI 1.33, 34.08), (Adjusted OR 11.27; 95% CI 1.83, 69.52).

**Maternal age:** Women who were greater than or equal to 25 years old were more likely to stay quit than women who were less than 25 years old in univariate analysis (Unadjusted OR: 8.92, 95% CI 1.35, 59.03). This association did not remain statistically significant in multivariate analysis (Adjusted OR 3.92; 95% CI 0.27, 57).

### **Discussion**

We were able to identify marginally significant associations between annual household income during pregnancy and use of WIC during pregnancy with women who quit smoking during pregnancy. Significant risk factors for staying quit, identified in this preliminary analysis, included maternal education, maternal age and postpartum depressive symptoms. Because of the limited sample size that was available (68 women who quit smoking during pregnancy compared to 85 women who smoked during pregnancy), this study only had adequate power to detect Odds Ratios of 4.0 or higher for quitting smoking during pregnancy and 5.0 or higher for staying quit, 2 years postpartum (24 women who stayed quit 2 year postpartum compared to 34 women who relapsed into smoking 2 years postpartum). Further research is warranted to re-examine some additional well-known risk factors such as stressful life events, social support and people in household who smoke in conjunction with staying quit. This research could be useful in highlighting characteristics of women who quit smoking during pregnancy and

providing some exploratory insight into characteristics of women who managed to stay quit, 2 years postpartum in order to be able to account for those characteristics while implementing smoking cessation interventions and ensuring efficient distribution of limited resources.

## Introduction

**Smoking during Pregnancy:** Maternal smoking during pregnancy has been causally associated with a multitude of adverse health outcomes such as fetal growth restriction, premature rupture of the membranes, placenta previa, placental abruption, pre-term delivery, and low birth weight babies. Babies born to women who smoke during pregnancy have about a 30% odds of being born prematurely, are more likely to be born with low birth weight (less than 2500 grams or 5.5 pounds) - increasing their risk for illness or death, weight an average of 200 grams or less than infants born to women who do not smoke and are 1.4 to 3 times more likely to die of Sudden Infant Death Syndrome (SIDS). In addition, in utero exposure to cigarette smoke can have negative long-term effects on the growth, development, and behaviors of offspring.<sup>39</sup>

Hospital and NICU costs for infants associated with maternal smoking during pregnancy are also a huge economic burden. A 2002 study done on 1997 PRAMS data collected from 13 states estimated the prevalence of smoking during pregnancy at 18%. This study examined an association of smoking during pregnancy with infant admission to the Neonatal Intensive Care Unit (NICU) and found that estimated average cost per night of stay, in the hospital, for a non-NICU infant was \$748, whereas for a NICU infant, a night of stay in the NICU nursery led to an average cost of \$2500.<sup>1</sup>

Adding to the evidence of harmful effects of smoking is an association between smoking status and food insecurity. Those who live in smoking households are more likely to be food insecure. Not only does spending on cigarettes divert resources away from investing in healthy foods, but the lost productivity resulting from diseases caused by smoking can lower income and raise the likelihood of food insecurity.<sup>11</sup>

Another factor of interest is pregnancy intention. Women's behavior during pregnancy may be influenced by their attitude towards pregnancy. Women with intended pregnancies are somewhat more likely to quit smoking during pregnancy than similar women with unintended pregnancies, but this varies according to the mother's social and demographic characteristics.<sup>22</sup>

A study in Maine conducted with 10 years of PRAMS data (1988-1997) found smoking prevalence to remain high in the last 3 months of pregnancy even though there was an overall decline in smoking prevalence. One of the characteristics of women who smoked through pregnancy, identified by this study, was WIC participation. Over the 10-year study period, even though there was a slight dip in smoking prevalence, women participating in WIC had consistently higher prevalence of smoking in the last 3 months of pregnancy compared to those women who did not participate in WIC.<sup>12</sup>

Furthermore, one of the adverse outcomes that are linked to high prevalence of depression among pregnant women and mental and physical health problems during pregnancy is continuation of high-risk behaviors, such as cigarette smoking.<sup>2</sup>

According to 2004 PRAMS data from 26 states the prevalence of smoking during pregnancy is estimated at 13%, which is lower than the 1997 numbers but still higher than national estimates of 10% and still far above the Healthy People 2010 goal of 1%. Younger, less educated, non-Hispanic, white women and American Indian women are more likely to smoke during pregnancy compared to their older, more educated counterparts.<sup>10</sup>

**Staying Quit, postpartum:** The past 15 years have seen a steady decrease in the number of women who smoke while pregnant, partly because of an overall decline in smoking rates among all women of childbearing age and partly, due to interventions targeting women during the prenatal period<sup>34</sup>. Smoking cessation during pregnancy significantly reduces the risks of fetal death, low birth weight, and other complications of pregnancy.

Continued abstinence postpartum also reduces children's exposure to environmental tobacco smoke and its associated health risks. According to a recent study by the Harvard School of Public Health, subjects first exposed to second hand smoke (SHS) between the ages of 0-25 have higher odds of developing lung cancer than those who were exposed to SHS after the age of 25<sup>6</sup>. Exposure to second hand smoke is also an important contributor to the future risks of impaired pulmonary function<sup>38</sup>. Moreover, sustained abstinence postpartum reduces women's lifetime risk for smoking-related diseases<sup>27</sup>.

In spite of overall decline of smoking prevalence, relapse into smoking postpartum remains high. According to a presentation on "Risk Factors for Smoking Cessation Relapse After Pregnancy" at the 9<sup>th</sup> Annual Maternal & Child Health Epidemiology Workshop in 2003, 50% of the PRAMS respondents successfully quit smoking during pregnancy. Among those who successfully quit smoking during pregnancy, almost 40% of them relapsed into smoking 2-6 months postpartum.<sup>13</sup>

There is evidence that suggests that the strategies used by women who quit smoking during pregnancy differ from those used by non-pregnant quitters, and thus, the resumption of smoking postpartum represents the end of a temporary suspension of smoking rather than a failure to maintain a long-term behavior change.<sup>25</sup>

A recent study findings published by researchers at University of North Carolina at Chapel Hill state: Women who remained smoke-free postpartum were bolstered by strong social support, strong internal belief systems, strong beliefs in postpartum health benefits of not smoking, negative experiences with a return to smoking and concrete strategies for dealing with temptations. Women who relapsed postpartum were undermined by easy access to cigarettes, reliance on cigarettes to deal with stress, lack of financial resources, lack of resources for childbearing and, low self-esteem.<sup>34</sup>

Prior studies have consistently reported the prevalence of relapse into smoking, postpartum at 50% or higher<sup>13, 23, 24, 25, 27, 35</sup>. More needs to be done to encourage women who quit smoking pregnancy, to continue to remain smoke-free, postpartum.

### **Specific Aims**

The data used in this study comes from the 2004 Oregon Pregnancy Risk Assessment Monitoring System (PRAMS) and from its 2006 longitudinal follow-up survey – PRAMS-2 – administered in 2006. This study aims to:

- 1) Establish prevalence of smoking at four different time points: 3 months before pregnancy (T1), last 3 months of pregnancy (T2), 2-6 months postpartum (T3), and, 2 years postpartum (T4).
- 2) Establish prevalence of staying quit among those who women who quit smoking during pregnancy at two time points: 2-6 months postpartum (T3) and 2 years postpartum (T4).
- 3) Identify characteristics of women who quit smoking during pregnancy compared to those women who continued to smoke through pregnancy.



4) Preliminary exploration into possible characteristics of women who stayed quit, 2 years postpartum, compared to those women who quit smoking during pregnancy but relapsed into smoking, two years postpartum.

Smoking still remains a serious problem that affects the health and welfare of the general population not only through the main effects of smoking but also through the exposure to second hand smoke (SHS). Limited information exists regarding correlates of those women who manage to remain smoke-free, postpartum. Information provided by this study could prove useful in examining factors associated with quitting smoking during pregnancy and staying quit, 2 years postpartum in order to target groups of women who may need to receive more effective and timely smoking cessation interventions.

## Methods

### PRAMS

This project uses data collected through the 2004 Oregon Pregnancy Risk Assessment Monitoring System (PRAMS) and its longitudinal follow-up – PRAMS-2 – administered in 2006. PRAMS is an epidemiologic surveillance program maintained by the Centers for Disease Control and Prevention (CDC) and implemented by participating state health departments. Research has indicated that maternal behaviors during pregnancy may influence infant birth weight and mortality rates. The goal of the PRAMS project is to improve the health of mothers and infants by reducing adverse outcomes such as low birth weight, infant mortality and morbidity, and maternal morbidity. PRAMS combines two modes of data collection; a survey conducted by mailed questionnaire with multiple follow-up attempts, and a survey by telephone. Here is the sequence of contacts for PRAMS surveillance:

- **Preletter.** This letter introduces PRAMS to the mother and informs her that a questionnaire will soon arrive.
- **Initial Mail Questionnaire Packet.** This packet is sent to all sampled mothers 3 to 7 days after the preletter. Its contents are described below.
- **Tickler.** The tickler serves as a thank you and a reminder note. It is sent 7 to 10 days after the initial mail packet.
- **Second Mail Questionnaire Packet.** This packet is sent to all sampled mothers who have not yet responded 7 to 14 days after the tickler has been sent.
- **Third Mail Questionnaire Packet.** This third packet is sent to all remaining nonrespondents 7 to 14 days after the second questionnaire.
- **Telephone Follow-up.** Telephone follow-up is initiated for all mail nonrespondents 7 to 14 days after mailing the last questionnaire.

The 2004 Oregon PRAMS survey was designed as an eighty-item self-administered questionnaire. Both the written survey and the interview ask the same questions. Oregon

PRAMS questionnaires and interviews were completed in either English or Spanish. A complete copy of the 2004 Oregon PRAMS survey is included in Appendix A.

## **PRAMS-2**

In January 2006 the Oregon Office of Family Health PRAMS team began re-surveying PRAMS respondents whose children had turned 24 months old. This new survey includes questions on health insurance, chronic diseases, oral health, well child-care, medical home, breastfeeding, smoking, domestic violence, family planning, child nutrition, immunization, early intervention, childcare, and reading to child.

PRAMS-2 was administered to all mothers who responded to the PRAMS Survey, with the exception of:

- Those who indicated “Do not contact me again” on the contact/information card included with PRAMS survey.
- Those whose babies were deceased.

From Jan06 to Aug06, the PRAMS-2 survey was only administered by mail. From Sep06 to present, the PRAMS-2 survey is administered by both mail and phone.

PRAMS-2 mailing materials are sent in English language to mothers who returned the PRAMS survey in English, and in Spanish language to mothers who returned the PRAMS survey in Spanish.

A complete copy of the 2006 Oregon PRAMS-2 survey is included in Appendix B. The PRAMS-2 survey format is similar to the PRAMS format with up to two mailings of the PRAMS-2 surveys to each mother and telephone follow-up with mothers who do not respond by mail.

PRAMS employs several complex weighting mechanisms in order to adjust for aspects of the subject selection methods. Because PRAMS oversamples for maternal race and ethnicity, a sampling weight is applied to the data. In the 2004 Oregon PRAMS, the

sampling weight was calculated to also account for oversampling based on low birth weight. The 2006 longitudinal follow-up, PRAMS-2, was re-weighted to account for loss to follow-up of the PRAMS respondents.

### **PRAMS/PRAMS-2 Weighting Methodology**

The population of interest for PRAMS is all mothers who are residents of Oregon who delivered within Oregon a live-born infant during the surveillance period. For PRAMS surveillance, there is often a particular interest from a public health perspective in certain subpopulations. These subpopulations may not represent a large portion of a state's overall population. To make inferences about specific subpopulations and make comparisons among several subpopulations, infants in those subpopulations (commonly called *strata*) will need to be oversampled (i.e., sampled at a higher rate than other subpopulations). The main advantage of *stratified sampling* is that it permits separate estimates of subgroups of interest and permits comparisons across these subgroups. Oregon chose to stratify by race/ethnicity (of mother) as defined by CDC: African American, American Indian/Alaskan Native, Asian/Pacific Islander, Hispanic, and White. This will enable Oregon to obtain accurate data about small minority race/ethnicity groups. Oversampling these groups should provide robust estimates of overall responses. In addition to the race/ethnicity stratification, a low birth weight white stratum was added, thus making the total number of strata for Oregon, six. Because of the recent reported increase in Oregon infant mortality deaths due to low birth weight and based on Oregon's birth population composition, it is feasible to over sample low birth weight babies only from the "White" strata.

Participation in Oregon PRAMS may vary based on additional maternal characteristics. That is, some women who share common traits may be less likely to respond to the survey than others. Because of this, a nonresponse weight is used to compensate for non-participation. Nonresponse weights are typically based on common demographic characteristics within each sampling stratum that are identified after the data has been collected and non-respondents have been compared to respondents.

Finally, participation in the Oregon PRAMS can also be affected by characteristics of the sampling scheme. That is, some women may be less likely than others to participate because they were not covered by the sampling frame. This situation can occur when there are accidental duplications in the birth certificate records. Duplicate records can lead to missing files in the group of selected participants. To adjust for women who may have been omitted in this fashion, a non-coverage weight is calculated and applied to the data. Factors that are related to non-coverage are identified by the CDC once all of the PRAMS data has been collected.

The longitudinal follow-up – PRAMS-2 – used the same three classification of weights that PRAMS uses, i.e., weighting for maternal race/ethnicity, non-response and, non-coverage. Only the non-response weight was re-calculated to account for those women that were lost to follow-up. All those women who also indicated “Do not contact me again” on the PRAMS information card were also included in the non-response weight category. For a detailed description of PRAMS weighting methodology, please refer to <http://www.cdc.gov/prams/methodology.htm>.

## Variable Coding

### Outcome

Only those women who reported smoking at least 100 cigarettes in the past 2 years were included in this analysis. There are two outcomes being examined in this analysis: (i) Quitting smoking during pregnancy – will be referred to as ‘Quit’ from now on, and, (ii) Staying quit, 2 years postpartum – will be referred to as ‘Staying quit’ from now on. All women who smoked 3 months before pregnancy and quit smoking during pregnancy were included in the quit outcome variable (Table 1). All those women who smoked 3 months before pregnancy, quit smoking during pregnancy and stayed quit, 2 years postpartum were included in the staying quit outcome variable (Table 2). For crosstab analysis, the “event” was coded as 1, else 0.

**Table 1: Coding of quitting smoking during pregnancy outcome variable (Among those who reported smoking at least 100 cigarettes in the past 2 years)**

Presence of Smoking (Yes/No)		Number of Women	
3 months before becoming pregnant	Last 3 months of pregnancy		
Yes	Yes	85	Smoked during pregnancy
Yes	No	68	Quit smoking during pregnancy

(Data source: 2004 PRAMS/2006 PRAMS-2)

**Table 2: Patterns of smoking among women who reported smoking at least 100 cigarettes in the past 2 years.**

Presence of Smoking (Yes/No)				Number of Women	
3 months before pregnancy (T1)	Last 3 months of pregnancy (T2)	2-4 months postpartum (T3)	2 years postpartum (T4)		
Yes	Yes	Yes	Yes	69	Persistent Smokers
Yes	Yes	Yes	No	5	Excluded
Yes	Yes	No	Yes	2	Excluded
Yes	Yes	No	No	5	Excluded
Yes	No	Yes	Yes	15	Quit/Relapsed
Yes	No	Yes	No	6	Excluded
Yes	No	No	Yes	19	Quit/Relapsed
Yes	No	No	No	24	Stayed Quit
No	Yes	Yes	Yes	0	Excluded
No	Yes	Yes	No	0	Excluded
No	Yes	No	Yes	0	Excluded
No	Yes	No	No	0	Excluded
No	No	Yes	Yes	3	Excluded
No	No	Yes	No	0	Excluded
No	No	No	Yes	2	Excluded
No	No	No	No	3	Excluded
Reported not smoking 100 cigarettes in the past 2 years				689	Excluded
Smoking information missing				23	Excluded
<b>Total</b>				<b>865</b>	

(Data source: 2004 PRAMS/2006 PRAMS-2)

### Independent Variables

The 2004 Oregon PRAMS assessed respondents for numerous experiences and behaviors related to pregnancy and birth outcomes. The 2004 PRAMS was also linked to birth certificates in order to access maternal demographic characteristics such as race/ethnicity, maternal education and maternal age. The 2006 longitudinal follow up assessed respondents' behavioral outcomes, 2 years postpartum, and outcomes related to

their now two-year old child. This analysis used information from both the PRAMS and PRAMS-2 surveys to attempt to profile women who quit smoking during pregnancy and explore potential correlates of women who stayed quit 2 years postpartum.

### **For the outcome of Quitting smoking during pregnancy (Quit)**

#### ***Variables Derived from Birth Certificate Information***

Variables drawn from birth certificate information used in this study, for analysis of quitting smoking during pregnancy, included maternal age, maternal race/ethnicity, education and marital status

Maternal race/ethnicity was separated into five categories: Hispanic, non-Hispanic White, non-Hispanic African American, non-Hispanic Asian/Pacific Islander, and non-Hispanic American Indian/Alaska Native. To ensure sufficient sample size in each category and because of the race/ethnicity distribution in Oregon<sup>40</sup>, the maternal race/ethnicity variable was dichotomized into Whites and non-Whites.

Information regarding maternal age (in years) was originally available as a continuous variable in the dataset. However, initial analysis revealed a nonlinear trend between age and the log odds of women quitting smoking during pregnancy. To improve the ease of analysis, this information was re-coded as a categorical variable. To ensure sufficient sample size in each category, maternal age was dichotomized into <25 years and  $\geq 25$  years.

Maternal education for the PRAMS variable analysis was coded as a categorical variable with 2 categories to be consistent with the maternal education variable in the PRAMS-2 dataset. The categories are <12<sup>th</sup> grade/12<sup>th</sup> grade or GED and >12<sup>th</sup> grade.



Table 3 includes a complete list of birth certificate variables used in this analysis, along with their original response options and recoded categories.

**Table 3. Independent variables derived from birth certificate  
(Used for the outcome of quitting smoking during pregnancy)**

<b>Birth Certificate Measure</b>	<b>Possible Responses</b>	<b>Coding for Analysis</b>
Maternal Age	Continuous	0 = <25 years 1 = 25 years or greater
Marital Status	Married Divorced/Separated/ Widowed	1 = All Else 2 = Married
Maternal Race/Ethnicity	- White - Hispanic - African American - Asian/Pacific Islander - American Indian/ Alaskan Native	0 = Whites 1 = non-Whites
Maternal Education	- Less than 12 <sup>th</sup> grade - 12 <sup>th</sup> grade - Greater than 12 <sup>th</sup> grade	0 = <12 <sup>th</sup> /12 <sup>th</sup> grade 1 = Greater than 12 <sup>th</sup> grade

(Linked with 2004 Oregon PRAMS data)

**For the outcome of Quitting smoking during pregnancy (Quit)**

***Variables derived from Oregon PRAMS***

The variables measured on the 2004 Oregon PRAMS used in this analysis, for the outcome of quitting smoking during pregnancy, included food insecurity during pregnancy, physician counseling for smoking during pregnancy, annual household income during pregnancy, pre-pregnancy smoking intensity, maternal age, intimate partner violence (IPV), stressful life events (SLE) during pregnancy, pregnancy intention, depressive symptoms during pregnancy and physical activity assessed 2-6 months postpartum.

According to a study by Frank Chaloupka, smoking households tend to experience higher levels of food insecurity <sup>11</sup>. Food insecurity during pregnancy was

measured using the question, “During the 12 months before your new baby was born, did you ever eat less than you felt you should because there wasn’t enough money to buy food?” Possible responses are “No” or “Yes”.

Some prior studies have assessed physical counseling for smoking during pregnancy<sup>9,14</sup>. Physician counseling for smoking during pregnancy was assessed with the question, “During any of your prenatal care visits, did a doctor, nurse or other health care worker talk with you about – How smoking during pregnancy could affect my baby (Please count only discussion, not reading materials or videos)” Possible responses were “No” or “Yes”.

Annual household income during pregnancy was divided into categories based upon percentage of the federal poverty threshold (% FPL), which is published by the Department of Health and Human Services. Because the 2004 Oregon PRAMS asks respondents about income in the year prior to giving birth, percentages of the FPL were based upon guidelines for 2003. All participants who reported an annual income were classified as earning either 0%—99 % FPL, 100%—184% FPL, or 185% FPL and greater. Cutoff points for each category were based upon their significance to public programs. In particular the Oregon WIC program requires that participants have incomes less than 185% FPL. The Oregon Food Stamp program now has an extended categorical eligibility which allows Oregon households to qualify for food stamps if they have incomes of upto 185% of FPL<sup>44</sup>. To account for the limited sample size in this study and since both WIC and the food stamps program use less than 185% FPL as the eligibility cut-off, this variable was further dichotomized into 0-184% FPL and  $\geq$ 185% FPL.

Pre-pregnancy smoking intensity was measured with the question, “In the 3 months before you got pregnant, how many cigarettes did you smoke on an average day?” The question also included information on how many cigarettes constitute a pack – “(A pack has 20 cigarettes).” Smokers were classified as “light” or “heavy” based on the number of cigarettes smoked on an average day. Those who smoked  $\leq 10$  cigarettes a day were classified as light smokers and those who smoked  $> 10$  cigarettes a day were classified as heavy smokers <sup>26</sup>.

Intimate Partner Violence (IPV) during pregnancy was measured using 2 items on the PRAMS questionnaire. They are, “During your most recent pregnancy, did an ex-husband or ex-partner push, hit, slap, kick, choke, or physically hurt you in any other way” – and – “During your most recent pregnancy, were you physically hurt in any way by your husband or partner?” Possible responses to either question were “No” or “Yes”. Respondents who answered “Yes” to either question were classified as having experienced IPV and only those who answered “No” to both questions were classified as not having experienced IPV during pregnancy.

Stressful life events have been examined as a risk factor associated with returning to smoking postpartum <sup>9</sup>, so this factor should also warrant examination in association with quitting smoking during pregnancy. Stressful life events were measured using a series of questions that assessed thirteen events that may have occurred during the 12 months prior to giving birth. Some of the events included on the survey were serious illness of a close family member, separation or divorce, homelessness, increased arguments with a husband or partner, and the loss of a job by either the mother or her husband or partner. All of the stressful life event measures included “Yes” or “No” as

possible responses. For this analysis, because of the limited sample size, these events were dichotomized into those women experiencing ‘0 events’ and those women experiencing 1 or more events in the year preceding delivery. Table 4 describes all the Stressful life events measured in PRAMS.

**Table 4. Stressful Life Event Measures, 2004 Oregon PRAMS  
(Used for the outcome of quitting smoking during pregnancy)**

Event
A close family member was very sick and had to go to the hospital
I got separated or divorced from my husband or partner
I moved to a new address
I was homeless
My husband or partner lost his job
I lost my job even though I wanted to go on working
I argued with my husband or partner more than usual
My husband or partner said that he didn’t want me to be pregnant
I had a lot of bills I couldn’t pay
I was in a physical fight
My husband or partner or I went to jail
Someone very close to me a bad problem with drinking or drugs
Someone very close to me died

Based on data from the 1988 NMIHS (National Maternal and Infant Health Survey) and and 1988 NSFG (National Survey of Family Growth), women’s behavior during pregnancy may be influenced by whether the pregnancy was planned or unplanned<sup>22</sup>. Pregnancy intention was measured using responses to the question, “Thinking back to just before you got pregnant with your new baby, how did you feel about becoming pregnant?” Women who reported either “I wanted to be pregnant sooner” or “I wanted to be pregnant then” were considered to have intended pregnancies. Women who reported

either “I wanted to be pregnant later” or “I didn’t want to be pregnant then or at any time in the future” were considered to have unintended pregnancies.

Depressive symptoms during pregnancy were measured using two items on the PRAMS questionnaire. The first question asked, “While you were pregnant, how often did you feel down, depressed, or hopeless?” The second question asked, “While you were pregnant, how often did you have little interest or pleasure in doing things?” Possible responses for both questions included “Always,” “Often,” “Sometimes,” “Rarely,” or “Never.” For this analysis, answers of “Always” or “Often” were considered a positive report of depressive symptoms. Answers of “Sometimes,” “Rarely,” and “Never” were considered a negative report of depressive symptoms. All respondents who provided a positive answer to either one or both of the questions were considered to have experienced depressive symptoms during pregnancy. All respondents who provided negative answers to both of the questions were considered to have been free of depressive symptoms during pregnancy. Table 5 includes a complete list of PRAMS variables used in this analysis (for the outcome of quitting smoking during pregnancy), along with their original response options and recoded categories.

**Table 5. (Independent variables derived from 2004 Oregon PRAMS  
(Used for the outcome of quitting smoking during pregnancy))**

<b>PRAMS Measure</b>	<b>Possible Responses</b>	<b>Coding for Analysis</b>
Food Insecurity during pregnancy	- No - Yes	1 = No 2 = Yes
Physician counseling for smoking during prenatal care visits	- No - Yes	0 = Yes 1 = No
Annual Household Income— during pregnancy	-Less than \$10,000 -\$10,000—\$14,999 -\$15,000—\$19,999 -\$20,000—\$24,999 -\$25,000—\$34,999 -\$35,000—\$49,999 -\$50,000 or more	0 = 0%—184% FPL 1 = 185% FPL or greater

Pre-Pregnancy Smoking Intensity	- 41 cigarettes or more - 21 to 40 cigarettes - 11 to 20 cigarettes - 6 to 10 cigarettes - 1 to 5 cigarettes - Less than 1 cigarette - None (0 cigarette)	1 = Heavy (>10 cigarettes) 2 = Light (<= 10 cigarettes)
Intimate Partner Violence— During Pregnancy, Current Husband or Partner/Ex- husband or Ex-Partner	-No -Yes	1 = Yes 2 = No
Stressful Life Events during pregnancy (13 events)	-No -Yes	1 = 1 Event or greater 2 = 0 Events
Pregnancy Intention	-I wanted to be pregnant sooner -I wanted to be pregnant later -I wanted to be pregnant then -I didn't want to be pregnant then or at any time in the future	1 = Unintended 2 = Intended
Depressive Symptoms during Pregnancy -Depressed Mood - Little Interest or pleasure	-Always -Often -Sometimes -Rarely -Never	Always/Often = Yes Sometimes/Rarely/Never = No
WIC Participation	-No -Yes	0 = Yes 1 = No

**For the outcome of staying quit 2 years postpartum (Staying Quit)**

***Variables derived from Oregon PRAMS***

Since, the outcome of staying quit assesses smoking status among PRAMS-2 respondents, 2 years after delivery; most variables used in this analysis were from PRAMS-2. But, there were a few variables from PRAMS that were meaningful to examine for inclusion in this analysis. Those variables include current physical activity and antenatal depressive symptoms, assessed 2-6 months postpartum.

Current physical activity was assessed with the following question, “In the past month, how many days a week did you get at least 30 minutes of physical activity of exercise? (For example, walking, dancing, yard work or sweeping).” Possible response choices included “Less than 1 day per week”, “1 to 4 days per week” and “5 or more days per week.” To account for the limited sample size, this variable was dichotomized into  $\leq 4$  days per week and  $\geq 5$  days per week. Table 4 includes a complete list of PRAMS variables used in this study, along with their original response options and recoded categories.

Antenatal depressive symptoms were assessed with two items on the PRAMS questionnaire. The first question asked, “Since your new baby was born, how often did you feel down, depressed, or hopeless?” The second question asked, “Since your new baby was born, how often did you have little interest or pleasure in doing things?” Possible responses for both questions included “Always,” “Often,” “Sometimes,” “Rarely,” or “Never.” For this analysis, answers of “Always” or “Often” were considered a positive report of depressive symptoms. Answers of “Sometimes,” “Rarely,” and “Never” were considered a negative report of depressive symptoms. All respondents who provided a positive answer to either one or both of the questions were considered to have experienced depressive symptoms after delivery. All respondents who provided negative answers to both of the questions were considered to have been free of depressive symptoms 2-6 months after delivery. Table 6 includes the list of PRAMS variables used in this analysis (for the outcome of staying quit), along with their original response options and recoded categories.

**Table 6. Independent variables derived from 2004 Oregon PRAMS  
(Used for the outcome of staying quit, 2 years postpartum)**

Physical Activity in the past month	- Less than 1 day per week - 1 to 4 days per week - 5 or more days per week	1 = 5 or more days per week 2 = 4 or less days per week
Antenatal Depressive Symptoms	- Always - Often	Always/Often = Yes Sometimes/Rarely/Never = No
- Depressed Mood	- Sometimes	
- Little Interest	- Rarely - Never	

**For the outcome of staying quit, 2 years postpartum (Staying Quit)**

***Variables derived from Oregon PRAMS-2***

The variables measured on the 2006 Oregon PRAMS-2 used in this analysis included food insecurity (13-24 months after delivery), people in the household who smoke, social support, marital status, maternal education, postpartum stressful life events (SLE), depressive symptoms in the first 12 months after delivery, depressive symptoms 13-24 months after delivery, current annual household income, maternal age, maternal physical activity and smoking cessation counseling from physician or health care worker (HCW).

Armour et al. found that families with income near the federal poverty level spend a large share of their income on cigarettes. This suggests an interaction between food insecurity and household income levels on smoking behavior<sup>5</sup>. With our limited sample size (n=58), we were unable to examine any interactions and so income and food insecurity were examined as individual risk factors for staying quit. Food Insecurity in PRAMS-2 was measured with the question, “In the past 12 months, did you ever eat less than you felt you should because there wasn’t enough money to buy food?” Possible responses to this question were “No” or “Yes”.



People in the household who smoked were measured with the question, “Not including yourself, is there anyone in your household who smokes cigarettes, cigars, or pipes?”

Possible responses are “No” or “Yes”.

Carmichael et al. cited the lack of information on social support as a limitation in their study examining correlates of postpartum smoking relapse<sup>9</sup>. Social support was measured using a series of questions which measured five events listed in Table 7. All of the social support questions had possible responses of “Yes” or “No”. Questions 2-5 were considered more relevant to the outcome of this analysis as an indicator of emotional social support and so only those questions were used in this analysis. To account for the limited sample size of this study, these questions were collapsed and dichotomized. Those who answered “No” to all four of the questions were considered as not having any social support and those who answered “Yes” to any one or more of the 4 questions were considered as having social support.

**Table 7. Social Support Measures, 2006 Oregon PRAMS-2**  
**Measures**

---

You have someone who would loan you money for food or bills if you needed it **(not used for purposes of this analysis)**.

---

You have someone who would help you if you were sick and needed to be in bed.

---

You have someone who would take you to the clinic or doctor’s office if you needed a ride.

---

You have someone you can count on to listen to you when you need to talk.

---

You have someone who shows you love and affection.

---

The PRAMS-2 survey includes questions regarding demographics such as marital status, maternal education and maternal age, assessed 2 year postpartum. These variables

were used for the analyses of staying quit rather than demographics from the birth certificate.

Marital status was measured with the question, “What is your current marital status?” Possible responses were “Never married”, “Married”, “Widowed”, “Divorced”, “Separated”. This variable was dichotomized into those who were married and all else to ascertain effect of being married on the outcome of interest.

Maternal education was measured with the question, “What is the highest level of school you have completed?” Possible responses were “Less than 12<sup>th</sup> grade”, “12<sup>th</sup> grade or GED” and “More than 12<sup>th</sup> grade”. This variable was dichotomized into all those who had a high school education or less and all those who had greater than a high school education.

Maternal age was dichotomized based on the most commonly used cut-off point among prior literature reviews for ease of interpretation. Thus, maternal age was dichotomized into <25 years and ≥25 years.

Stressful life events in the year preceding delivery were found to be significantly associated with return to smoking, 2-6 months postpartum<sup>9</sup>. Hence, stressful life events between 13-24 months after delivery would warrant examination in association with staying quit, 2 years postpartum. Stressful life events were measured using a series of questions that assessed thirteen events that may have occurred during the past 12 months. It is the same series of questions that is reported in Table 3. This variable was dichotomized, because of the small sample size and cell counts, into those who experienced no SLEs’ - “None” and those who experienced at least one SLE – “≥1 event”.

Allen et al. , using 2004 PRAMS data from 16 states, found that smokers with depressive symptoms are more likely to relapse after attempting to quit than those without depressive symptoms<sup>3</sup>. Maternal depressive symptoms 2 years after delivery were measured using 2 items on the PRAMS-2 questionnaire. The first item examines depressive symptoms in the first 12 months after delivery with the question, “During the FIRST 12 months of your two-year-old’s life, was there a period of two or more weeks when almost every day you: (a) Felt sad, blue or depressed for most of the day, (b) Lost interest or pleasure in most things that you usually cared about or enjoyed.” Possible responses for both (a) and (b) are “No” or “Yes”. The second item examines depressive symptoms in the past 12 months with the question, “In the PAST 12 months, has there been a period of two or more weeks when almost every day you: (a) Felt sad, blue or depressed for most of the day, (b) Lost interest or pleasure in most things you usually cared about or enjoyed.” Possible responses for both (a) and (b) are “No” or “Yes”. If the respondent answered “Yes” to either part (a) or (b) for the first 12 months, then they were classified as having depressive symptoms during that time period. If they answered “No” to both parts (a) and (b), then they were classified as not having depressive symptoms during that time period. Assessing depressive symptoms in the respondents during the past 12 months was done similarly.

Current annual household income was divided into categories based upon percentage of the federal poverty threshold (% FPL). All participants who reported an annual income were classified as earning either 0%—184 % FPL or  $\geq$ 185%.

Levine et al. published an article assessing concerns about weight and its association with smoking behavior in women<sup>24</sup>. Current physical activity was assessed

using the question, “In the past month, how many days a week did you get at least 30 minutes of physical activity or exercise? (For example, walking, dancing, yard work or sweeping).” Possible responses were “Less than 1 day per week”, “1 to 4 days per week” and “5 or more days per week”. To account for the small sample size, this variable was further dichotomized into  $\leq 4$  days per week and  $\geq 5$  days per week.

Postpartum smoking cessation counseling by physician or HCW was assessed with the following two questions – “During any of your healthcare visits in the last 12 months, did a doctor, nurse, or other health care worker talk with you about any of the things listed below? (i) Advise you to quit smoking, (ii) Offer you help on how to quit smoking” Possible responses were “Yes” or “No”. Both these items were examined separately for the outcome of interest. Table 8 includes a complete list of PRAMS-2 variables used in this analysis, along with their original response options and recoded categories.

**Table 8. Independent variables derived from 2006 Oregon PRAMS-2  
(Used for the outcome of staying quit, 2 years postpartum)**

<b>PRAMS-2 Measure</b>	<b>Possible Responses</b>	<b>Coding for Analysis</b>
Food Insecurity in the past 12 months	- No - Yes	1 = No 2 = Yes
HCW advise to quit smoking	- No - Yes	0 = Yes 1 = No
HCW offer help to quit smoking	- No - Yes	0 = Yes 1 = No
Current Annual Household Income—	-Less than \$10,000 -\$10,000—\$14,999 -\$15,000—\$19,999 -\$20,000—\$24,999 -\$25,000—\$34,999 -\$35,000—\$49,999 -\$50,000 or more	1 = 0%—184% FPL 2 = 185% FPL and greater
People in household who smoke	- No - Yes	1 = No 2 = Yes
Social Support (4 out of 5 questions)	- No - Yes	1 = No 2 = Yes
Marital Status	- Never Married	0 = Married

	- Married - Widowed - Divorced - Separated	1 = All else
Maternal Education	- Less than 12 <sup>th</sup> grade - 12 <sup>th</sup> grade or GED - More than 12 <sup>th</sup> grade	0 = Less than 12 <sup>th</sup> /12 <sup>th</sup> 1 = More than 12 <sup>th</sup> grade
Stressful Life Events (13 events)	-No -Yes	0 = 0 Events 1 = 1 or more events
Postpartum depressive symptoms in first 12 months	- No - Yes	0 = No 1 = Yes
Postpartum depressive symptoms in past 12 months	- No - Yes	0 = No 1 = Yes
Maternal Age	- Continuous	0 = <25 years 1 = 25 years or greater
Current physical Activity	- Less than 1 day per week - 1 to 4 days per week - 5 or more days per week	1 = 5 or more days per week 2 = 1 to 4 days per week

### **Data Management**

Tasks related to cleaning and editing of PRAMS data files, including the correction of errors and inconsistencies, are the responsibility of state health departments. For the 2004 Oregon PRAMS and the 2006 Oregon PRAMS-2, the Department of Human Services (DHS) performed all data entry verification and telephone interview monitoring (as all telephone interviews were performed by a hired contractor). The data files were then checked for consistency through an automated process at the CDC. Finally, the CDC created the Oregon PRAMS analysis file, complete with analysis weights. This file was then provided to Oregon DHS. Re-weighting for the PRAMS-2 data set was done at DHS with instructions from the CDC.

For this project, I acquired the PRAMS merged with the PRAMS-2 follow up data file in STATA format. All analyses for this project were performed using STATA Version 10 (STATA Corporation) software package. Because Oregon PRAMS and

PRAMS-2 responses are de-identified and do not contain personal identifying information, the Institutional Review Board (IRB) of Oregon Health and Science University exempted this project from review.

## **Statistical Analysis**

### Descriptive Analysis

Weighted data was used for all parts of the statistical analysis. The prevalence of smoking at the four different time points was examined. The prevalence of staying quit, 2 years postpartum, was also determined among the PRAMS-2 respondents. Cross tabulations were then generated to determine the number of women who quit smoking during pregnancy and the number of women who stayed quit, within each of the covariates. These cross tabulations were examined to determine if there were sufficient cell counts within each of the variable categories to be eligible for inclusion in a multivariable model.

### Univariate Analysis

Simple logistic regression models were constructed to determine unadjusted odds ratios (ORs) to examine significant risk factors between those women who quit smoking during pregnancy compared to those who continued to smoke during pregnancy. The same was done for the women who stayed quit, 2 years postpartum, compared to those who relapsed into smoking, 2 years postpartum.

All correlates with a p-value of  $\leq 0.25$  were eligible for inclusion in the multivariable logistic regression model. A large number of prior studies that have assessed smoking behaviors during pregnancy and 2-6 months postpartum have included demographics in their analyses<sup>3, 9, 13, 21, 26, 27, 28, 43</sup> and so, demographics such as maternal

age, maternal education, marital status, annual household income (% FPL) and maternal race/ethnicity were included in the model, to adjust for, even if the p-value was greater than 0.25.

### Multivariate analysis

Multivariate analysis was conducted using Hosmer & Lemeshow's applied logistic regression technique for model building<sup>46</sup>. Expected counts for each variable were examined to ensure adequate sample sizes in each group. Some of the variables were further collapsed into dichotomized formats if cell sizes were too small to obtain accurate inferences. Continuous variables such as maternal age were categorized if the relationship between the log odds of the outcome and associated covariate showed departure from linearity. All examined independent covariates with sufficient cell counts and with a p-value of  $\leq 0.25$  was eligible for inclusion into the multivariable logistic regression model.

### Forward stepwise model building

STATA does not allow for automated model selection processes such as backwards, forwards, stepwise processes or best subset selection using weighted data. Hence, forward manual stepwise was performed by adding the variables to the model by the level of significance in univariate analysis. The variable that was most significant was added first to the model and additional variables added, one at a time, based on the significance of the association with the outcome of quitting smoking during pregnancy.

Similarly, a forward manual stepwise model building process was performed for the outcome of staying quit, 2 years postpartum.

## Results

### Summary

#### *Response Rates*

The 2004 Oregon PRAMS sampled 1,968 women. The weighted response rate for PRAMS was 74.8%. The PRAMS-2 survey was sent to 1,935 of these 1,968 women. 865 women responded to the PRAMS-2 survey with an un-weighted response rate of 44%. The weighted response rate, or more appropriately, the percent of the Oregon population that this represents, is 51.1%. The overall combined response rate for both PRAMS and PRAMS-2 was 38%. The weighted response rate for PRAMS-2 was calculated using the formula defined by the CDC. For detailed information on calculation of the PRAMS-2 response rate and the overall response rate, please refer to Appendix C.

#### *Demographics of PRAMS-2 respondents*

All of the percentages reported here are weighted. The majority of the PRAMS-2 respondents in this study were greater than or equal to 25 years old (78.3%), had more than a high school education (62.6%) and were married (76.4%). Annual household income was split right down the middle with half the respondents earning an annual income of less than 185% of the federal poverty level (49.4%) and the other half earning an annual income of greater than or equal to 185% of the federal poverty level (50.6%). Most respondents also reported not smoking at least 100 cigarettes in their entire life (64.6%) and were non-Hispanic Whites (71%)

#### *Smoking Prevalence*



20.9% of the PRAMS-2 respondents smoked at least 100 cigarettes in the past two years (N=167). The PRAMS survey was able to establish smoking prevalence during three time points – 3 months before pregnancy (T1), last 3 months of pregnancy (T2) and 6 months postpartum (T3) Information about smoking 3 months before pregnancy was available for 853 respondents. Those who reported smoking less than 100 cigarettes in the past 2 years, were coded as non-smoking for each of those time points. 19% of all PRAMS-2 respondents reported smoking less than one or more cigarettes on an average day in the 3 months before pregnancy (N=154). Information about smoking in the last 3 months of pregnancy was available for 854 respondents. In the last 3 months of pregnancy, 11% of the respondents (N=87) smoked less than one or more cigarettes on an average day. Information about smoking 6 months postpartum was available for 855 respondents. At 6 months postpartum, 14% of the respondents (N=104) reported smoking less than one or more cigarettes on an average day.

35.4% of the PRAMS-2 respondents smoked at least 100 cigarettes in their entire life (N=284). The PRAMS-2 survey was able to establish smoking prevalence at a fourth time point (T4) – 2 years postpartum. Information about smoking 2 years postpartum was available for 853 respondents. At 2 years postpartum, 15% of the respondents (N=135) smoked less than one or more cigarettes on an average day.

#### *Persistent Smokers*

Those PRAMS-2 respondents who reported smoking less than one or more cigarettes on an average day at all the four time points – 3 months before pregnancy (T1), last 3 months of pregnancy (T2), 6 months postpartum (T3) and 2 years postpartum (T4) – were classified as persistent smokers. Smoking information at all the four time points

were available for 830 respondents. 8.7% of the respondents (N=69) reported smoking at all the four time points.

#### *Stayed Quit*

For purposes of this analysis, those respondents who reported smoking before pregnancy (T1), quit smoking during pregnancy (T2) and continued to remain smoke-free at 6 months postpartum (T3) and 2 years postpartum (T4) were classified as staying quit. 3.7% of the respondents (N=24) were classified as staying quit.

#### *Relapsers*

Those respondents who reported smoking before pregnancy (T1), quit during pregnancy (T2) and relapsed into smoking, 2 years postpartum (T4) were classified as relapsers. 2.6% of the respondents (N=34) were classified as relapsers.

#### *Quit smoking during pregnancy*

Among those PRAMS-2 respondents who reported smoking 3 months before pregnancy (T1), prevalence of quitting smoking during pregnancy was 44.5% (N=85).

#### *Stayed quit, 2-6 months postpartum*

Among those PRAMS-2 respondents who reported smoking 3 months before pregnancy (T1) and reported quitting smoking during pregnancy (T2), prevalence of staying quit 2-6 months postpartum (T3) was 65% (N=47).


#### *Stayed quit, 2 years postpartum*

Among those PRAMS-2 respondents who reported smoking 3 months before pregnancy (T1) and reported quitting smoking during pregnancy (T2), prevalence of those who stayed quit 2-6 months postpartum (T3) and continued to stay quit 2 years postpartum (T4) was 59% (N=24).

*Non-Smokers*

Those respondents who did not smoke at any of the four time points were classified as non-smokers. This category also included those who reported not smoking at least 100 cigarettes in the past 2 years or in their entire life. 82% of the respondents (703) were classified as non-smokers. Figure 1 is a flowchart of the four time points under consideration.

**Figure 1**



**Outcome: Quit smoking during pregnancy**

68 women reported quitting smoking during pregnancy while 85 women reported smoking during pregnancy with a 44.5% prevalence of quitting smoking during pregnancy.

*Maternal Education*

Information regarding maternal education was available for 83 women who smoked during pregnancy and all 68 women who quit smoking during pregnancy. 40% of those who had a high school education or less (38) and 58% of those who had greater than a high school education (30) quit smoking during pregnancy.

*Marital Status*

Information regarding maternal marital status was available for all 85 women who smoked during pregnancy and all 68 women who quit smoking during pregnancy. 52% of those who were married (36) and 39% of those who were not married (includes never married, divorced and separated) (32) quit smoking during pregnancy.

#### *Maternal Age*

Information regarding maternal age was available for all 85 women who smoking during pregnancy and all 68 women who quit smoking during pregnancy. 38% of women aged less than 25 years (33) and 55% of women 25 years old or greater (35) quit smoking during pregnancy.

#### *Maternal race/ethnicity*

Information regarding maternal race/ethnicity was available for all 85 women who smoked during pregnancy and all 68 women who quit smoking during pregnancy. 45% of non-Hispanic Whites (37) and 44% of non-Whites (includes American Indian/Alaskan Native, African American, Asian/Pacific Islander and Hispanic) (31) quit smoking during pregnancy.

#### *Food insecurity during pregnancy*

Information regarding food insecurity during pregnancy was available for all 85 women who smoked during pregnancy and 67 women who quit smoking during pregnancy. 42% of those who didn't report food insecurity (52) and 50% of those who reported food insecurity (15) quit smoking during pregnancy.

#### *HCW talk about smoking during prenatal visits*

Information regarding doctor, nurse or health care worker (HCW) talking about smoking was available for 84 women who smoked during pregnancy and 67 women who

quit smoking during pregnancy. 40% of those who reported that a doctor, nurse or other HCW spoke with them about how smoking during pregnancy could affect the baby (57) and 66% of those who didn't report any HCW talking about smoking, quit smoking during pregnancy (10).

*Annual household income during pregnancy (% FPL)*

Information regarding annual pre-pregnancy household income was available for 80 women who smoked during pregnancy and 65 women who quit smoking during pregnancy. 37% of those who had annual household income during pregnancy of 0-184% FPL (37) and 68% of those who had annual household income during pregnancy of  $\geq$ 185% FPL (28) quit smoking during pregnancy.

*Pre-pregnancy smoking intensity*

Information regarding pre-pregnancy smoking intensity was available for 85 women who smoked during pregnancy and 68 women who quit smoking during pregnancy. 35% of heavy smokers (20) and 52% of light smokers (48) quit smoking during pregnancy.

*Intimate Partner Violence (IPV) during pregnancy*

Information regarding IPV was available for all 85 women who smoked during pregnancy and all 68 women who quit smoking during pregnancy. 37% of those who reported experiencing IPV during pregnancy (7) and 46% of those who reported not experiencing any IPV during pregnancy (61) and quit smoking during pregnancy.

*Stressful Life Events (SLE) during pregnancy*

Information regarding SLE during pregnancy was available for all 85 women who smoked during pregnancy and all 68 women who quit smoking during pregnancy. 43%

of those who reported experiencing 1 or more SLE's during pregnancy (54) and 56% of those who reported experiencing no SLE's (14) quit smoking during pregnancy.

#### *Pregnancy Intention*

Information regarding pregnancy intention was available for 82 women who smoked during pregnancy and 67 women who quit smoking during pregnancy. 40% of those who reported unintended pregnancies (35) and 54% of those who reported intended pregnancies (32) quit smoking during pregnancy.

#### *Depressive symptoms during pregnancy*

Information regarding depressive symptoms during pregnancy was available for 83 women who smoked during pregnancy and 67 women who quit smoking during pregnancy. 47% of women who reported experiencing depressive symptoms during pregnancy (5) and 44% of women who reported experiencing no depressive symptoms during pregnancy (63) quit smoking during pregnancy.

#### *WIC during pregnancy*

Information regarding WIC use during pregnancy was available for all 85 women who smoked during pregnancy and all 68 women who quit smoking during pregnancy. 37% of women who reported using WIC during pregnancy (39) and 63% of women who reported not using WIC during pregnancy (29), quit smoking during pregnancy.

#### *Pre-pregnancy BMI*

Information regarding pre-pregnancy weight was available for 81 women who smoked during pregnancy and all 68 women who quit smoking during pregnancy. 58% of women who were considered normal (40) based on BMI classification and 36% of

those who were classified as all else (included underweight, obese and overweight) (28), quit smoking during pregnancy.

#### *Other variables*

Other variables that were assessed but were not included for analysis due to insufficient cell counts were alcohol consumption during pregnancy.

#### **Univariate Analysis**

The proportion of women, who quit smoking during pregnancy, did not vary by most of the correlates examined in this analysis. In univariate logistic analyses annual household income during pregnancy and WIC participation during pregnancy were of marginal significance as risk factors for women quitting smoking during pregnancy.

Table 9 describes the associations between women who quit smoking during pregnancy and possible correlates, including Unadjusted ORs and 95% CIs.

Women who quit smoking during pregnancy were more likely to have higher annual household incomes during pregnancy (OR: 3.68, 95% CI: 0.98, 13.79) and were also more likely to have not used WIC during pregnancy compared to women who smoked during pregnancy (OR: 2.84, 95% CI: 0.84, 9.21).

**Table 9: Risk Factors for Quitting smoking during pregnancy (n=68)**  
 Compared to women who smoked through pregnancy (n=85)

Maternal characteristic	Numerator (un-weighted)	Denominator (un-weighted)	Quit (weighted)	Unadjusted OR (95% CI)
<b>Total</b>	68	153	44.5%	
<b>Maternal Education</b>				
<12 <sup>th</sup> /12 <sup>th</sup> grade	38	99	40%	Referent 2.10 (0.70, 6.30)
>12 <sup>th</sup> grade	30	52	58%	
Missing		2		
<b>Marital Status</b>				
All else	32	84	39%	Referent 1.68 (0.59, 4.82)
Married	36	69	52%	
<b>Maternal Age</b>				
<25 years	33	81	38%	Referent 2.02 (0.70, 5.80)
>= 25 years	35	72	55%	
<b>Maternal Race/Ethnicity</b>				
Non-Whites	31	68	44%	Referent 1.02 (0.39, 2.66)
Non-Hispanic Whites	37	85	45%	
<b>Food insecurity during pregnancy</b>				
No	52	115	42%	Referent 1.39 (0.45, 4.25)
Yes	15	37	50%	
Missing	1	1		
<b>HCW talk about smoking during prenatal visits</b>				
Yes	57	132	40%	Referent 2.85 (0.68, 11.93)
No	10	19	66%	
Missing	1	2		
<b>Annual Household Income during Pregnancy</b>				
0-184% FPL	37	101	37%	Referent 3.68 (0.98, 13.79)
>= 185% FPL	28	44	68%	
Missing	3	8		
<b>Pre-pregnancy Smoking Intensity</b>				
Heavy	20	64	35%	Referent 2.08 (0.71, 6.06)
Light	48	89	52%	
<b>Intimate Partner Violence during pregnancy</b>				
Yes	7	26	37%	Referent 1.47 (0.33, 6.48)
No	61	127	46%	
<b>Stressful Life Events during pregnancy</b>				
>=1 Event(s)	54	132	43%	Referent 1.70 (0.37, 7.77)
0 Events	14	21	56%	
<b>Pregnancy Intention</b>				
Unintended	35	86	40%	Referent 1.77 (0.61, 5.16)
Intended	32	63	54%	
Missing	1	4		
<b>Depressive Symptoms</b>				



<b>during Pregnancy</b>				
Yes	22	55	47%	Referent 0.90 (0.29, 2.78)
No	45	95	44%	
Missing	1	3		
<b>WIC during pregnancy</b>				
Yes	39	104	37%	Referent 2.84 (0.87, 9.21)
No	29	49	63%	
<b>Maternal Pre-pregnancy BMI</b>				
All Else	28	74	36%	Referent 2.49 (0.84, 7.40)
Normal (18.5<= BMI < 25.0)	40	75	58%	
Missing		4		

(Data source: 2004 PRAMS/2006 PRAMS-2)

### **Multivariate Logistic Analysis**

For the outcome of quitting smoking during pregnancy, all covariates that were had a p-value of  $\leq 0.25$  were included in the multivariable model. Demographic factors such as maternal race/ethnicity, maternal age and maternal education were included in the model to adjust for.

The final model for women who quit smoking during pregnancy included the variables of maternal education, marital status, maternal age, maternal race/ethnicity, HCW talk about smoking during prenatal visit, pre-pregnancy smoking intensity, annual household income during pregnancy, the use of WIC during pregnancy and pre-pregnancy BMI. Table 10 describes the associations from the multivariable model between women who quit smoking during pregnancy and possible correlates, including Adjusted ORs and 95% CIs.

**Table 10: Multivariable Model - Risk Factors for Quitting smoking during pregnancy (n=68)**

Compared to women who smoked through pregnancy (n=85)

Maternal characteristic	Numerator (un-weighted)	Denominator (un-weighted)	Quit (weighted)	Unadjusted OR (95% CI)	Adjusted OR (95% CI)
<b>Total</b>	68	153	44.5%		
<b>Maternal Education</b>					
<12 <sup>th</sup> /12 <sup>th</sup> grade	38	99	40%	Referent 2.10 (0.70, 6.30)	Referent 1.64 (0.35, 7.66)
>12 <sup>th</sup> grade	30	52	58%		
Missing		2			
<b>Marital Status</b>					
All else	32	84	39%	Referent 1.68 (0.59, 4.82)	Referent 0.89 (0.25, 3.14)
Married	36	69	52%		
<b>Maternal Age</b>					
<25 years	33	81	38%	Referent 2.02 (0.70, 5.80)	Referent 1.60 (0.44, 5.76)
>= 25 years	35	72	55%		
<b>Maternal Race/Ethnicity</b>					
Non-Whites	31	68	44%	Referent 1.02 (0.39, 2.66)	Referent 0.82 (0.25, 2.72)
Non-Hispanic Whites	37	85	45%		
<b>Food insecurity during pregnancy</b>					
No	52	115	42%	Referent 1.39 (0.45, 4.25)	Excluded from model
Yes	15	37	50%		
Missing	1	1			
<b>HCW talk about smoking during prenatal visits</b>					
Yes	57	132	40%	Referent 2.85 (0.68, 11.93)	Referent 2.31 (0.41, 12.83)
No	10	19	66%		
Missing	1	2			
<b>Annual Household Income during Pregnancy</b>					
0-184% FPL	37	101	37%	Referent 3.68 (0.98, 13.79)	Referent 2.17 (0.50, 9.42)
>= 185% FPL	28	44	68%		
Missing	3	8			
<b>Pre-pregnancy Smoking Intensity</b>					
Heavy	20	64	35%	Referent 2.08 (0.71, 6.06)	Referent 1.84 (0.54, 6.31)
Light	48	89	52%		
<b>Intimate Partner Violence during pregnancy</b>					
Yes	7	26	37%	Referent 1.47 (0.33, 6.48)	Excluded from model
No	61	127	46%		
<b>Stressful Life Events during pregnancy</b>					
>=1 Event(s)	54	132	43%	Referent 1.70 (0.37, 7.77)	Excluded from model
0 Events	14	21	56%		
<b>Pregnancy Intention</b>					
Unintended	35	86	40%	Referent 1.77 (0.61, 5.16)	Excluded from model
Intended	32	63	54%		
Missing	1	4			

<b>Depressive Symptoms during Pregnancy</b>					
Yes	22	55	47%	Referent 0.90 (0.29, 2.78)	Excluded from model
No	45	95	44%		
Missing	1	3			
<b>WIC during pregnancy</b>					
Yes	39	104	37%	Referent 2.84 (0.87, 9.21)	Referent 1.08 (0.29, 4.03)
No	29	49	63%		
<b>Maternal Pre-pregnancy BMI</b>					
All Else	28	74	36%	Referent 2.49 (0.84, 7.40)	Referent 1.71 (0.54, 5.35)
Normal (18.5 <= BMI < 25)	40	75	58%		
Missing		4			

(Data source: 2004 PRAMS/2006 PRAMS-2)

### **Forward Manual Stepwise Model Building**

The variable that was first added to the manual stepwise model was annual household income during pregnancy with the highest level of significance in univariate analysis ( $p=0.05$ ). The next variable to be added was use of WIC during pregnancy ( $p=0.08$ ). Even though there was moderate correlation between household income and WIC participation, inclusion or exclusion of either variable did not change the overall results and significance of the remaining variables in the final model. Pre-pregnancy BMI, HCW talk about smoking, maternal education, pre-pregnancy smoking intensity, maternal age, marital status and maternal race/ethnicity were added in the order listed. Table 11 includes results of manual stepwise selection and its corresponding ORs and p-values.

**Table 11. Forward Manual Stepwise - Multivariable Logistic Regression Model, Odds Ratios and p-values for quitting smoking during pregnancy**

Characteristic	Stage 1 OR, (p-value)	Stage 2 OR, (p-value)	Stage 3 OR, (p-value)	Stage 4 OR, (p-value)	Stage 5 OR, (p-value)	Stage 6 OR, (p-value)	Stage 7 OR, (p-value)	Stage 8 OR, (p-value)	Stage 9 OR, (p-value)
Annual household income during pregnancy	3.68 (0.05)	2.98 (0.15)	2.44 (0.21)	2.17 (0.26)	2.14 (0.28)	2.18 (0.27)	2.08 (0.30)	2.13 (0.31)	<b>2.17 (0.30)</b>
WIC during pregnancy		1.64 (0.48)	1.34 (0.64)	1.40 (0.60)	1.22 (0.75)	1.14 (0.84)	1.06 (0.92)	1.07 (0.92)	<b>1.08 (0.91)</b>
Pre-pregnancy BMI			1.93 (0.27)	1.93 (0.25)	1.72 (0.34)	1.63 (0.40)	1.69 (0.36)	1.69 (0.36)	<b>1.70 (0.36)</b>
HCW talk about smoking				2.28 (0.30)	2.40 (0.30)	2.44 (0.30)	2.23 (0.35)	2.24 (0.34)	<b>2.31 (0.34)</b>
Maternal education					1.67 (0.45)	1.98 (0.34)	1.66 (0.51)	1.66 (0.51)	<b>1.64 (0.53)</b>
Pre-pregnancy smoking intensity						1.92 (0.29)	1.86 (0.32)	1.86 (0.31)	<b>1.84 (0.33)</b>
Maternal age							1.56 (0.49)	1.61 (0.46)	<b>1.60 (0.47)</b>
Marital status								0.90 (0.87)	<b>0.89 (0.86)</b>
Maternal race/ethnicity									<b>0.82 (0.74)</b>

(Data source: 2004 PRAMS/2006 PRAMS-2)

### **Outcome: Stayed Quit, 2 years postpartum**

Among the 68 women who quit smoking during pregnancy, 6 were excluded because they reported smoking at 2-6 months postpartum. 4 women had missing information regarding smoking status at 2 years postpartum. Among the remaining 58 women, 24 stayed quit at 2 years postpartum while 34 relapsed into smoking 2 years postpartum.

#### *Maternal Education*

Information regarding maternal education was available for all 24 who stayed quit and 34 relapsers. 14.6% of those who had a high school education or less (7) and 77.7% of those who had more than a high school education stayed quit, 2 years postpartum (17).

#### *Marital Status*

Information regarding marital status was available for all 24 who stayed quit and all 34 relapsers. 60.3% of those who were married (17) and 54.1% of those who were not married (includes never married, divorced and separated) (7) stayed quit, 2 years postpartum.

*Maternal race/ethnicity*

Information regarding maternal race/ethnicity was available for all 24 who stayed quit and all 34 relapsers. 62.1% of non-Hispanic Whites (12) and 40.1% of non-Whites (12) stayed quit, 2 years postpartum.

*Current annual household income (% FPL)*

Information regarding current annual household income was available for all 24 who stayed quit and all 34 relapsers. 53.8% of those who were in the 0-184% FPL (11) and 63.8% of those who were in the  $\geq 185\%$  FPL (13) stayed quit, 2 years postpartum.

*Maternal age*

Information regarding maternal age was available for all 24 who stayed quit and all 34 relapsers. 53.7% of those who were less than 31 years old (15) and 72.3% of those who were greater than or equal to 31 years old (9) stayed quit, 2 years postpartum.

*Food Insecurity in the past 12 months*

Information regarding food insecurity in the past 12 months was available for all 24 who stayed quit and all 34 relapsers. 57.3% of those who reported experiencing food insecurity in the past 12 months (5) and 64.3% of those who reported experiencing no food insecurity (19) stayed quit, 2 years postpartum.

*People in household who smoke*

Information regarding people in household who smoke was available for all 24 who stayed quit and all 34 relapsers. 63.3% of those who reported not having any one in the household that smokes (17) and 50.1% of those who reported having someone in the household who smokes (7) stayed quit, 2 years postpartum.

*Pre-pregnancy smoking intensity*

Information regarding pre-pregnancy smoking intensity was available for all 24 who stayed quit and all 34 relapsers. 55.3% of heavy smokers (7) and 60.9% of light smokers (17) stayed quit, 2 years postpartum.

*Stressful Life Events (SLEs) in the past 12 months*

Information regarding SLEs in the past 12 months was available for all 24 who stayed quit and all 34 relapsers. 37.3% of those who experienced 1 or more SLEs in the past 12 months (18) and 62.7% of those who experienced no SLEs in the past 12 months stayed quit, 2 years postpartum (6).

*Depressive symptoms in the first 12 months after delivery*

Information regarding depressive symptoms in the first 12 months after delivery was available for all 24 who stayed quit and all 34 relapsers. 49.7% of those who reported experiencing no depressive symptoms for 2 or more weeks in the first 12 months after delivery (13) and 63.5% of those who reported experiencing depressive symptoms for 2 or more weeks in the first 12 months after delivery (10) stayed quit, 2 years postpartum.

*Depressive symptoms 13-24 months after delivery*

Information regarding depressive symptoms 13-24 months after delivery was available for 22 who stayed quit and all 34 relapsers. 46.1% of those who reported

experiencing depressive symptoms for 2 or more weeks in the past 12 months (15) and 84% of those who reported experiencing depressive symptoms almost every day for a period of two or more weeks in the past 12 months (7) stayed quit, 2 years postpartum.

*Physical activity in the past month (at 6 months postpartum)*

Information regarding physical activity in the past month (at 6 months postpartum) was available for 23 who stayed quit and 32 relapsers. 33.3% of those who reported having 30 minutes of physical activity, in the last month, for 5 or more days in a week (4) and 44.2% of those who reported having 30 minutes of physical activity, in the last month, for 4 or less days in a week (19) stayed quit, 2 years postpartum.

*Health Care Worker (HCW) advice to quit smoking*

Information regarding HCW advice to quit smoking was available for 21 who stayed quit and 33 relapsers. 36.6% of those who received advice on quitting smoking from a HCW in the last 12 months (7) and 68% of those who did not receive advice on quitting smoking from a HCW in the last 12 months (14) stayed quit, 2 years postpartum.

*Health Care Worker (HCW) offer help to quit smoking*

Information regarding HCW offer help to quit smoking was available for 22 who stayed quit and 32 relapsers. 41.1% of those who were offered help to quit smoking by a HCW in the last 12 months (7) and 67.1% of those who were not offered help to quit smoking

by a HCW in the last 12 months (15) stayed quit, 2 years postpartum.

*Current BMI*

Information regarding current BMI was available for all 24 who stayed quit and for all 34 relapsers. 62% of those who were normal (8) according to BMI classification

and 57% of those who were underweight, overweight or obese (16) stayed quit, 2 years postpartum.

### *Other Variables*

Other variables that were examined but not included for analysis due to insufficient sample sizes were antenatal depressive symptoms, assessed 2-6 months postpartum, social support, physical activity in the past month at 2 years postpartum and alcohol consumption in an average week in the past 12 months (assessed 2 years postpartum).

### **Univariate Analysis**

In simple logistic analyses maternal education, maternal age and depressive symptoms 13-24 months after delivery were significantly associated with women staying quit, 2 years postpartum.

Women who quit smoking during pregnancy and continued to stay quit, 2 years postpartum were more likely to have greater than a high school education (Unadjusted OR: 20.33, 95% CI: 4, 103.54) and were more likely to be 25 years old or greater (Unadjusted OR: 8.92, 95% CI: 1.35, 59.03). Women who stayed quit were also more likely to report experiencing depressive symptoms in the second year following pregnancy (Unadjusted OR: 6.15, 95% CI: 1.15, 32.79).

Table 12 describes the associations between women who stayed quit and possible correlates, including Unadjusted ORs and 95% CIs.



**Table 12: Risk Factors for Staying Quit, 2 years postpartum (n=24)**

Compared to women who relapsed into smoking, 2 years postpartum (n=34)

<b>Maternal characteristic</b>	<b>Numerator (un-weighted)</b>	<b>Denominator (un-weighted)</b>	<b>Stayed Quit (weighted)</b>	<b>Unadjusted OR (95% CI)</b>
<b>Total</b>	24	58	59%	
<b>Maternal Education</b>				
<12 <sup>th</sup> /12 <sup>th</sup> grade	7	25	15%	Referent 20.33 (3.99, 103.54)
>12 <sup>th</sup> grade	17	33	78%	
<b>Marital Status</b>				
All Else	7	21	54%	Referent 1.29 (0.18, 9.05)
Married	17	37	60%	
<b>Maternal Race/Ethnicity</b>				
non-Whites	12	27	40%	Referent 2.44 (0.49, 12.19)
Whites	12	31	62%	
<b>Annual Household Income</b>				
0-184% FPL	11	33	54%	Referent 1.51 (0.25, 9.22)
>= 185% FPL	13	25	64%	
<b>Maternal Age</b>				
<25 years	15	37	54%	Referent 8.92 (1.35, 59.03)
>=25 years	9	20	72%	
Missing		1		
<b>Food Insecurity (past 12 months)</b>				
Yes	5	11	57%	Referent 1.34 (0.12, 14.57)
No	19	47	64%	
<b>People in household who smoke</b>				
Yes	7	23	50%	Referent 1.72 (0.27, 11.07)
No	17	35	63%	
<b>Pre-pregnancy Smoking Intensity</b>				
Heavy (>10 cigs)	7	18	55%	Referent 1.25 (0.2, 7.8)
Light (<=10 cigs)	17	40	61%	
<b>Stressful Life Events in the past 12 months</b>				
>=1 Event	18	46	37%	Referent 3.03 (0.34, 27.07)
0 Events	6	12	63%	
<b>Depressive symptoms first year after delivery</b>				
No	13	35	50%	Referent 1.76 (0.27, 11.30)
Yes	10	22	64%	
Missing	1	1		
<b>Depressive symptoms 2nd year after delivery</b>				
No	15	41	46%	Referent 6.73 (1.33, 34.08)
Yes	8	16	85%	
Missing	1	1		
<b>Physical Activity (6 months postpartum)</b>				
>=5 days a week	4	12	33%	Referent 1.98 (0.17, 23.15)
<= 4 days a week	19	43	44%	

Missing	1	3		
<b>HCW advised to quit smoking</b>				
Yes	7	33	37%	Referent 3.68 (0.49, 27.89)
No	14	21	68%	
Missing	3	4		
<b>HCW offer help to quit smoking</b>				
Yes	7	24	41%	Referent 2.92 (0.40, 21.4)
No	15	30	67%	
Missing	2	4		
<b>Current BMI</b>				
All Else	16	36	57%	Referent 1.24 (0.20, 7.70)
Normal	8	22	62%	

(Data source: 2004 PRAMS/2006 PRAMS-2)

### **Multivariate Analysis**

For the outcome of staying quit 2 years postpartum, all covariates that had a p-value of  $\leq 0.25$  were included in the multivariable logistic regression model.

Demographic factors such as maternal race/ethnicity, maternal age, maternal education and annual household income (% FPL) were included in the model to adjust for.

The final model for staying quit included the variables of maternal education, marital status, maternal age, maternal race/ethnicity, depressive symptoms 13-24 months after delivery and HCW advice to quit smoking. Maternal education and annual household income were moderately correlated and annual household income was not significant in univariate analysis, hence, annual household income was dropped from the final model to avoid collinearity. Table 13 describes the multivariate associations between women who stayed quit, 2 years postpartum, and the covariates, including Adjusted ORs and 95% CIs.

**Table 13: Multivariable Model - Risk Factors for Staying Quit, 2 years postpartum (n=24)**

Compared to women who relapsed into smoking, 2 years postpartum (n=34)

Maternal characteristic	Numerator (un-weighted)	Denominator (un-weighted)	Stayed Quit (weighted)	Unadjusted OR (95% CI)	Adjusted OR (95% CI)
<b>Total</b>	24	58	58.7%		
<b>Maternal Education</b>					
<12 <sup>th</sup> /12 <sup>th</sup> grade	7	25	14.6%	Referent	Referent
>12 <sup>th</sup> grade	17	33	77.7%	20.33 (3.99, 103.54)	13.30 (2.36, 74.87)
<b>Marital Status</b>					
All Else	7	21	54.1%	Referent	Referent
Married	17	37	60.3%	1.29 (0.18, 9.05)	0.35 (0.02, 5.04)
<b>Maternal Race/Ethnicity</b>					
non-Whites	12	27	40.1%	Referent	Referent
Whites	12	31	62.1%	2.44 (0.49, 12.19)	3.00 (0.23, 38.34)
<b>Annual Household Income</b>					
0-184% FPL	11	33	53.8%	Referent	Excluded because of collinearity with maternal education
>= 185% FPL	13	25	63.8%	1.51 (0.25, 9.22)	
<b>Maternal Age</b>					
<25 years	4	16	30%	Referent	Referent
>=25 years	20	41	80%	8.92 (1.35, 59.03)	3.92 (0.27, 57)
Missing		1			
<b>Food Insecurity (past 12 months)</b>					
Yes	5	11	57.3%	Referent	Excluded from model
No	19	47	64.3%	1.34 (0.12, 14.57)	
<b>People in household who smoke</b>					
No	17	35	63.3%	Referent	Excluded from model
Yes	7	23	50.1%	1.72 (0.27, 11.07)	
<b>Pre-pregnancy Smoking Intensity</b>					
Heavy (>10 cigs)	7	18	55.3%	Referent	Excluded from model
Light (<=10 cigs)	17	40	60.9%	1.25 (0.2, 7.8)	
<b>Stressful Life Events in the past 12 months</b>					
>=1 Event	18	46	37.3%	Referent	Excluded from model
0 Events	6	12	62.7%	3.03 (0.34, 27.07)	
<b>Depressive Symptoms (first 12 months after delivery)</b>					
No	13	35	49.7%	Referent	Excluded from model
Yes	10	22	63.5%	1.76 (0.27, 11.30)	
Missing	1	1			
<b>Depressive Symptoms (13-24 months after</b>					

<b>delivery)</b>					
No	15	41	46.1%	Referent	Referent
Yes	7	15	84%	6.73 (1.33, 34.08)	11.27 (1.83, 69.52)
Missing	2	2			
<b>Physical Activity (6 months postpartum)</b>					
>=5 days a week	4	12	33.3%	Referent	Excluded from model
<= 4 days a week	19	43	44.2%	1.98 (0.17, 23.15)	
Missing	1	3			
<b>HCW advised to quit smoking</b>					
Yes	7	33	36.6%	Referent	Referent
No	14	21	68%	3.68 (0.49, 27.89)	6.53 (0.99, 43.01)
Missing	3	4			
<b>HCW offer help to quit smoking</b>					
Yes	7	24	41.1%	Referent	Excluded from the model
No	15	30	67.1%	2.92 (0.40, 21.4)	
Missing	2	4			
<b>Current BMI</b>					
All Else	16	36	57%	Referent	Excluded from the model
Normal	8	22	62%	1.24 (0.20, 7.70)	

(Data source: 2004 PRAMS/2006 PRAMS-2)

### **Forward Manual Stepwise Model Building**

The variable that was first added to the manual stepwise model was maternal education ( $p=0.0005$ ). Maternal age was next; addition of maternal age to a model that already contained the variable of maternal education rendered the age/staying quit association not statistically significant. Depressive symptoms in 13-24 months after delivery, HCW talk about quitting smoking, maternal race/ethnicity and marital status were added to the model in the order listed. The variables of maternal education and depressive symptoms 13-24 months after delivery remained significant after inclusion of all other covariates. Table 14 includes results of manual stepwise selection and corresponding ORs and 95% CIs.

**Table 14. Forward Manual Stepwise - Multivariable Logistic Regression Model, Odds Ratios and p-values**

Characteristic	Stage 1 OR, (p-value)	Stage 2 OR, (p-value)	Stage 3 OR, (p-value)	Stage 4 OR, (p-value)	Stage 5 OR, (p-value)	Stage 6 OR, (p-value)
Maternal Education	20.33 (0.0005)	11.46 (0.006)	15.25 (0.001)	14.49 (0.003)	12.02 (0.01)	<b>13.30</b> <b>(0.004)</b>
Maternal age		4.33 (0.17)	3.51 (0.25)	2.90 (0.37)	3.27 (0.33)	<b>3.92</b> <b>(0.31)</b>
Depressive symptoms in the past 12 months			8.39 (0.015)	17.86 (0.002)	18.29 (0.003)	<b>11.27</b> <b>(0.01)</b>
HCW talk about quitting smoking				5.94 (0.04)	5.63 (0.06)	<b>6.53</b> <b>(0.05)</b>
Maternal race/ethnicity					2.29 (0.49)	<b>2.99</b> <b>(0.39)</b>
Marital status						<b>0.35</b> <b>(0.43)</b>

(Data source: 2004 PRAMS/2006 PRAMS-2)

## Discussion

### Summary

Prevalence of quitting smoking during pregnancy among the PRAMS-2 respondents was 44.5%. Prior studies have found a range of quitting rates during pregnancy – from 35% to 56% - <sup>35, 23, 21</sup> because the definition of quitting smoking during pregnancy varied among the different studies. Prevalence of staying quit among PRAMS-2 respondents was 59%. This measure has been assessed, by prior studies, consistently at 5-6 months postpartum and staying quit rates at that time have been estimated to be about 50% or lower <sup>13, 23, 24, 25, 27, 35</sup>. This study is one of the first studies examining prevalence of staying quit, 2 years postpartum. This may be an over-estimation partly because a higher percentage of women, who were lost to follow-up reported smoking at 2-6 months postpartum, compared to PRAMS-2 respondents and also because women may under-report smoking due to the social stigma attached to it.

This study of Oregon women found that annual household income during pregnancy and using WIC during pregnancy were of marginal significance in conjunction with quitting smoking while pregnant. After adjusting for maternal age, maternal race/ethnicity, maternal marital status, pre-pregnancy BMI and HCW talk about smoking during pregnancy, women with higher household incomes and women who did not participate in WIC during pregnancy were more likely to quit smoking, but these associations did not remain statistically significant.

This study also found that maternal education, maternal age and self-reported maternal depressive symptoms 13-24 months after delivery were significantly associated with staying quit, 2 years postpartum. After adjusting for other factors such as maternal

marital status, maternal race/ethnicity and HCW advice to quit smoking, the associations between maternal education and self-reported maternal depressive symptoms 13-24 months after delivery with the outcome of staying quit still remained statistically significant.

#### Comparison with previous findings

##### ***Prevalence of smoking at all 4 time points: before pregnancy (T1), during pregnancy (T2), 6 months postpartum (T3) and 2 years postpartum (T4)***

10.5% of the PRAMS-2 respondents reported smoking at all the four time points. Since this was one of the first studies assessing smoking prevalence, 2 years postpartum and the first study using PRAMS-2 data, this finding cannot be validated at the current time for external consistency.

##### ***Prevalence of staying quit***

Among those who quit smoking during pregnancy, 58.7% stayed quit, 2 years postpartum, while 41.3% relapsed into smoking, postpartum. According to 1996 PRAMS data, prevalence of relapse into smoking, postpartum, was 50.9% at 2-6 months postpartum.<sup>9</sup> The prevalence of relapse at 2 years, which our study found, is lower than these estimates. Other studies have consistently measured prevalence of relapse into smoking, postpartum, at about 50%. Our estimates are close to that. Also, these estimates are from the longitudinal follow-up and those lost to follow-up in the PRAMS-2 may have been more likely to be smokers. Thus, this number may be an overestimation of the true prevalence of staying quit in the underlying population.

##### ***Risk Factors for quitting smoking during pregnancy***

The results of this study found that annual household income during pregnancy and the use of WIC during pregnancy were of marginal significance in association with

quitting smoking during pregnancy. Women who do not participate in WIC during pregnancy are more likely to quit smoking during pregnancy. This is consistent with prior studies <sup>12</sup>. This study also found positive associations between maternal education, marital status, maternal age, pre-pregnancy smoking intensity, IPV, SLEs during pregnancy and pregnancy intention as risk factors for quitting smoking during pregnancy, albeit, not statistically significant.

The association between food insecurity during pregnancy and quitting smoking during pregnancy differed from existing literature. Our study found that those who reported experiencing food insecurity during pregnancy were more likely to quit smoking during pregnancy compared to those who reported not experiencing food insecurity, though this was not statistically significant. Prior studies found that smoking households tend to experience higher levels of food insecurity <sup>11</sup>. Prior studies also suggest the possibility of food insecurity being an effect modifier in the association between household income levels and smoking status. Smoking prevalence seems to be higher among low income families who were food insecure than it was for low income families who were food secure <sup>5</sup>. Our data did not have enough sample size to investigate any possible effect modifiers. This could be one possible explanation for our counter-intuitive findings. Un-weighted numbers result in higher rates of quitting smoking among women who are food secure while weighted numbers flip the association around. This could also be due to the complex weighting mechanism used by PRAMS to represent the ethnic distribution of the population represented by PRAMS.



### ***Risk Factors for staying quit, 2 years postpartum***

This study found significant associations between the variables of maternal education and self-reported maternal depressive symptoms 13-24 months after delivery with women staying quit, 2 years postpartum. Our findings regarding maternal education is consistent with prior studies that measured postpartum smoking prevalence at 2-6 months postpartum.<sup>10, 14, 21, 41</sup> Women with greater than a high school education are more likely to stay quit than those with a high school education or less. There is very limited information available on staying quit, 2 years postpartum to further validate these results.

Our study also found that women who reported experiencing depressive symptoms 13-24 months after delivery are more likely to stay quit. A 2004 PRAMS study examined data from 16 states that had elected to include the optional question of postpartum depressive symptoms on the survey. This study found that women who reported experiencing postpartum depressive symptoms (2-6 months postpartum) were less likely to stay quit and more likely to relapse into smoking, postpartum<sup>3</sup>. Our findings are counter-intuitive to this study. These results may be due to the fact that women reporting depressive symptoms 13-24 months after delivery may be chronic depressives and not typical postpartum depressives. There have been studies done examining the use of anti-depressants as an aid for smoking cessation<sup>45</sup>, which may have been the cause of our findings. Also, this is preliminary exploration of the first year of PRAMS-2 data; combining a few years of data is needed to further stratify this purported association by factors such as race/ethnicity and annual household income. Prior research has also found that nicotine dependence and depression often occur together in the same patient. But, very little research has been done to find out whether depressed

patients who quit smoking attain long-lasting abstinence and what the effects of quitting are on their depressive symptoms. It is uncertain whether smokers with a history of depression run an increased risk of a new depressive episode once they quit smoking<sup>32</sup>.

### Strengths and Limitations

This analysis used data from the 2004 Oregon Pregnancy Risk Assessment Monitoring System and its two-year longitudinal follow-up, the 2006 Oregon PRAMS-2. The greatest strength of using PRAMS data is its population-based nature. A representative sample of the state's population is surveyed thus making the results generalizable to the women throughout the state. The PRAMS-2 data were further re-weighted, not only, using the same weighting scheme for PRAMS, but also to account for those women that were lost to follow-up. Thus, PRAMS-2 also enjoys the generalizability that PRAMS provides.

Both PRAMS and the PRAMS-2 surveys evaluate a variety of measures that could affect maternal smoking status before, during and after pregnancy, such as stressful life events, social support and depressive symptoms during pregnancy and postpartum. This allowed for examination of a multitude of factors, in multivariate analysis, that could possibly influence maternal smoking behavior.

A third strength is the first-time use of 2006 PRAMS-2 data. The 2006 Oregon PRAMS-2 dataset is a longitudinal survey following the 2004 PRAMS respondents in time. Thus temporality of data was maintained as well as being able to shed some light on the demographic distribution of the PRAMS-2 respondents.

This study also had several limitations, the foremost being the lost to follow-up. As evidenced by the response rate, a sizeable number of women who were lost to follow-

**Table 15: Characteristics of PRAMS-2 respondents versus those lost to follow-up**

Maternal Characteristic	PRAMS-2 respondents		Lost to follow-up	
	Un-weighted count (%)	Weighted %	Un-weighted count (%)	Weighted %
Reported smoking 100 cigarettes in the past 2 years	167 (19.5)	21	282 (27)	30
Smoking 3 months before pregnancy	154 (18)	19	259 (25)	28.5
Smoking in the last 3 months of pregnancy	87 (10.2)	10.2	148 (14)	17.4
Smoking 2-6 months postpartum	104 (12.2)	13.6	200 (19)	22.3
Pre-pregnancy smoking intensity (Heavy smokers)	64 (41.6)	47.3	95 (36.7)	50.4
BC Maternal marital status (Married)	633 (75.6)	72.4	620 (56.2)	57
Food insecurity during pregnancy (Yes)	93 (12.8)	13	149 (14.2)	14.5
IPV during pregnancy (Yes)	28 (3.5)	2.3	41 (4.4)	2.4
Unintended pregnancies	300 (35.2)	31.2	498 (45.9)	43.7
Depressive symptoms during pregnancy (Yes)	163(19.4)	14.1	311 (29.8)	25.9
Depressive symptoms 2-6 months postpartum (Yes)	46 (5.4)	4.2	88 (8.3)	7.3
Used WIC during pregnancy	351 (41.1)	34.8	643 (60.6)	58.2
Physical Activity, 6 months postpartum (<=4 days a week)	660 (77.9)	77.3	802 (76.1)	72.4
Annual household income during pregnancy (0-184% FPL)	359 (43.8)	38.2	611 (63.2)	57.4
Maternal race/ethnicity (Whites)	389 (45.1)	79.8	311 (28.2)	62.2
Maternal age (<25)	220 (25.4)	25.4	417 (46.9)	52.2

**(Data source: 2004 Oregon PRAMS)**

up with the PRAMS-2 survey. In examining the characteristics of the PRAMS-2 respondents compared to those lost to follow-up, some differences were found. A higher proportion of women reported smoking at least 100 cigarettes in the past 2 years among those lost to follow-up (30%) compared to the PRAMS-2 respondents (21%). Smoking prevalence at the 3 time points measured in PRAMS, i.e. 3 months before pregnancy (T1), last 3 months of pregnancy (T2) and 2-6 months postpartum (T3) were higher in those lost to follow-up than among the PRAMS-2 respondents. Among those lost to follow-up, more women participated in WIC during pregnancy (58.2%), had unintended pregnancies (43.7%), experienced depressive symptoms during pregnancy (25.9) and 2-6 months postpartum (7.3%) compared to the PRAMS-2 respondents. Finally, women lost to follow-up were more likely to have been of lower income (57.4% in 0-184% FPL) compared to the PRAMS-2 respondents (38.2% in 0-184% FPL) and more likely to be less than 25 years of age. But, as stated before, the PRAMS-2 data were re-weighted again to account for the loss to follow-up and so our results may not have been highly influenced by those lost to follow-up. Table 15 provides a list of maternal characteristics and weighted distributions of PRAMS-2 respondents in comparison with those lost to follow-up.

Yet another limitation is the cross-sectional nature of PRAMS. Only PRAMS variables were used for the outcome of quitting smoking during pregnancy. Thus one cannot infer causation from the results of this analysis. It would not be possible to state whether smoking during pregnancy causes one to use WIC or whether using WIC during pregnancy causes one to continue smoking during pregnancy.

In examining the outcome of staying quit at 2 years postpartum, variables from PRAMS were either not meaningful to use or were not used due to insufficient sample sizes, thus, this study lacked the ability to use the temporality of data to its advantage.

A third limitation of this study is the small sample sizes. Thus the study did not have adequate power to detect meaningful shifts nor did the study have the ability to examine any effect modification.

The fourth limitation is that smoking status was self-reported and was not verified by any biological tests (like cotinine levels).<sup>9</sup> This combined with the fact that maternal smoking during pregnancy/postpartum has a social stigma attached to it may lead to under-reporting of the actual smoking prevalence in the PRAMS/PRAMS-2 population. There is also a possibility of reporting bias of sensitive topics such as IPV and alcohol consumption during pregnancy, thus, under-estimating the true Odds Ratios.

## Public Health Implications

This study finding lends further credibility to the assertion that annual household income is associated with quitting smoking during pregnancy and that maternal education levels are associated with continued abstinence, postpartum.

### **For the outcome of quitting smoking during pregnancy:**

Among the PRAMS-2 respondents 25.9% of those eligible for WIC participation during pregnancy (had annual household income during pregnancy of 0-184% FPL) did not use WIC. Oregon is one of just a handful of states to implement a federal food stamp policy option that allows Oregon households to qualify for food stamps if they have incomes of up to 185% of the FPL. WIC also uses the 185% FPL as cut-off for eligibility. Encouraging more eligible women to use WIC through continued awareness would be advantageous. Because WIC is a prenatal nutrition and health education program serving low-income women and children, WIC can provide opportunities for intervention and follow-up of women who are pregnant and smoke <sup>11</sup>.

The participants of the Oregon Food Stamp program, WIC program or TANF would also have lower annual household incomes and this would aid in focusing the limited resources available to those group of women who need more help to quit smoking during pregnancy.

### **For the outcome of staying quit, 2 years postpartum**

Women with greater than a high school education were more likely to stay smoke-free, 2 years postpartum. Smoking cessation interventions for quitting during pregnancy and staying quit, postpartum, should be targeted more towards women with lower levels of education. Since education levels and annual household income exhibit a

positive correlation, this would also mean targeting those women who report lower annual household incomes, akin to quitting smoking during pregnancy.

Women who reported experiencing depressive symptoms 13-24 months after delivery were also more likely to stay quit, postpartum. Depression and smoking are co-morbidities and should not be treated in isolation from each other<sup>40</sup>. Nicotine withdrawal may produce depressive symptoms or precipitate a major depressive episode, and thus, there may be some anti-depressants that might also be used as an aid for smoking cessation<sup>45</sup>. There is very little research done on how postpartum depressive symptoms can affect smoking behaviors in new mothers, especially those with a history of depressive symptoms. It is not clear whether remaining abstinent can cause an episode of experiencing depressive symptoms in smokers with a history of depression<sup>32</sup>. Health care workers need to be aware of this co-morbidity and the fact that women who report depressive symptoms 13-24 months after live birth may be chronic depressives, not typical postpartum depressives. Continued screening of depressive symptoms would be important to learn more about this health condition and how it can affect smoking behavior in women.

## Future Studies

This study lacked the required sample size to study the effect of certain well known correlates such as pre-pregnancy smoking intensity, depressive symptoms during pregnancy, people in household who smoke, stressful life events during pregnancy and postpartum and social support, for the outcomes of quitting smoking during pregnancy or staying quit, 2 years postpartum. Combining multiple years of PRAMS-2 data will render adequate sample size to study the effect of these variables on quitting smoking and staying quit. Enhanced understanding of behavioral differences by race/ethnicity of the survey population will be necessary to apply population-based interventions.

In this study, non-Hispanic Whites were more likely to quit smoking during pregnancy and more likely to stay quit, 2 years postpartum, though not statistically significant. Since, the Oregon population is predominantly non-Hispanic White <sup>39</sup>, minority race/ethnicity categories such as African Americans make up small target populations that may be amenable to culturally sensitive smoking cessation interventions.

It would be worth examining whether those mothers who reported experiencing depressive symptoms 13-24 months after delivery were diagnosed as having depression and whether they were on any anti-depressants. It would also be very insightful to examine patterns of self-reported depressive symptoms at all the different time points that this information is available for: during pregnancy, 2-6 months postpartum, in the first 12 months after delivery and 13-24 months after delivery.



## Summary and Conclusion

This was the first and foremost study to explore year one of the Oregon PRAMS-2 dataset.

Exploring risk factors for quitting smoking during pregnancy, we found marginally significant associations between annual household income during pregnancy and WIC participation during pregnancy with the outcome of quitting smoking during pregnancy. Although similar associations were observed in multivariate analysis, they were not statistically significant.

In our preliminary analysis of staying quit, 2 years postpartum, women who had higher levels of education and those who reported experiencing depressive symptoms in 13-24 months after delivery were more likely to stay quit, both in univariate and multivariate analysis.

We lacked the capability of using the temporality of the longitudinal PRAMS-2 dataset to our advantage. Hence, the cross-sectional nature of our analysis prohibited us from determining direction and causality of the associations.

The findings of this study suggest that additional well-known risk factors be re-examined with multiple years' of data to validate current preliminary findings. There is also a need for improved understanding of self-reported maternal depressive symptoms in the first two years after delivery and its effect on smoking behaviors of mothers.

## **Appendix A**

### 2004 Oregon Pregnancy Risk Assessment Monitoring System Questionnaire

First, we would like to ask a few questions about you and the time before you got pregnant with your new baby. Please check the box next to your answer.

1. *Just before you got pregnant, did you have health insurance?* Do not count Oregon Health Plan or Medicaid.

- No  
 Yes

2. *Just before you got pregnant, were you on Oregon Health Plan or Medicaid?*

- No  
 Yes

3. *During the month before you got pregnant with your new baby, how many times a week did you take a multivitamin or a prenatal vitamin?* These are pills that contain many different vitamins and minerals.

- I didn't take a multivitamin or a prenatal vitamin at all  
 1 to 3 times a week  
 4 to 6 times a week  
 Every day of the week

4. What is *your* date of birth?

19  
 Month Day Year

5. *Just before you got pregnant with your new baby, how much did you weigh?*

Pounds OR  Kilos

6. How tall are you without shoes?

Feet  Inches

OR  Centimeters

7. *Before you got pregnant with your new baby, did you ever have any other babies who were born alive?*

- No  Go to Question 10  
 Yes

8. Did the baby born *just before your new one weigh 5 pounds, 8 ounces (2.5 kilos) or less at birth?*

- No  
 Yes

9. Was the baby *just before your new one born more than 3 weeks before its due date?*

- No  
 Yes

The next questions are about the time when you got pregnant with your *new* baby.

10. Thinking back to *just before you got pregnant with your new baby, how did you feel about becoming pregnant?*

Check one answer

- I wanted to be pregnant sooner  
 I wanted to be pregnant later  
 I wanted to be pregnant then  
 I didn't want to be pregnant then or at any time in the future

**11. When you got pregnant with your new baby, were you trying to get pregnant?**

No

Yes → **Go to Question 15**

**12. When you got pregnant with your new baby, were you or your husband or partner doing anything to keep from getting pregnant?**

(Some things people do to keep from getting pregnant include not having sex at certain times [rhythm] or withdrawal, and using birth control methods such as the pill, condoms, cervical ring, IUD, having their tubes tied, or their partner having a vasectomy.)

No

Yes → **Go to Question 14**

**13. What were your or your husband's or partner's reasons for not doing anything to keep from getting pregnant?**

**Check all that apply**

- I didn't mind if I got pregnant
- I thought I could not get pregnant at that time
- I had side effects from the birth control method I was using
- I had problems getting birth control when I needed it
- I thought my husband or partner or I was sterile (could not get pregnant at all)
- My husband or partner didn't want to use anything
- Other → Please tell us:

---

**If you or your husband or partner was not doing anything to keep from getting pregnant, go to Question 15.**

**14. When you got pregnant with your new baby, what were you or your husband or partner doing to keep from getting pregnant?**

**Check all that apply**

- Tubes tied or closed (female sterilization)
- Vasectomy (male sterilization)
- Pill
- Condoms
- Shot once a month (Lunelle®)
- Shot once every 3 months (Depo-Provera®)
- Contraceptive patch (OrthoEvra®)
- Diaphragm, cervical cap, or sponge
- Cervical ring (NuvaRing® or others)
- IUD (including Mirena®)
- Rhythm method or natural family planning
- Withdrawal (pulling out)
- Not having sex (abstinence)
- Other → Please tell us:

---

**The next questions are about the prenatal care you received during your most recent pregnancy. Prenatal care includes visits to a doctor, nurse, or other health care worker before your baby was born to get checkups and advice about pregnancy.** (It may help to look at the calendar when you answer these questions.)

**15. How many weeks or months pregnant were you when you were *sure* you were pregnant?** (For example, you had a pregnancy test or a doctor or nurse said you were pregnant.)

Weeks **OR**  Months

I don't remember

**16. How many weeks or months pregnant were you when you had your first visit for prenatal care?** Do not count a visit that was only for a pregnancy test or only for WIC (the Special Supplemental Nutrition Program for Women, Infants, and Children).

Weeks **OR**  Months

I didn't go for prenatal care

**17. Did you get prenatal care as early in your pregnancy as you wanted?**

- No  
 Yes  
 I didn't want prenatal care →

**Go to Page 4, Question 19**

**18. Here is a list of problems some women can have getting prenatal care.** For each item, circle **Y** (Yes) if it was a problem for you during your most recent pregnancy or circle **N** (No) if it was not a problem or did not apply to you.

- |                                                                                   | <b>No</b> | <b>Yes</b> |
|-----------------------------------------------------------------------------------|-----------|------------|
| a. I couldn't get an appointment when I wanted one . . . . .                      | N         | Y          |
| b. I didn't have enough money or insurance to pay for my visits . . . . .         | N         | Y          |
| c. I had no way to get to the clinic or doctor's office . . . . .                 | N         | Y          |
| d. I couldn't take time off from work . . .                                       | N         | Y          |
| e. The doctor or my health plan would not start care as early as I wanted . . . . | N         | Y          |
| f. I didn't have my Oregon Health Plan or Medicaid card . . . . .                 | N         | Y          |
| g. I had no one to take care of my children . . . . .                             | N         | Y          |
| h. I had too many other things going on . . . . .                                 | N         | Y          |
| i. I didn't want anyone to know I was pregnant . . . . .                          | N         | Y          |
| j. Other . . . . .                                                                | N         | Y          |
- Please tell us:

---

**If you did not go for prenatal care, go to Question 25.**

**19. Where did you go *most of the time* for your prenatal visits? Do not include visits for WIC.**

**Check one answer**

- Hospital clinic
- Health department clinic
- Private doctor's office or HMO clinic
- Midwife's office
- At home
- Other —————> Please tell us:

---

**20. How was your prenatal care paid for?**

**Check all that apply**

- Oregon Health Plan or Medicaid
- Personal income (cash, check, or credit card)
- Health insurance or HMO (including insurance from your work or your husband's work)
- Indian Health Service
- Other —————> Please tell us:

---

**21. During any of your prenatal care visits, did a doctor, nurse, or other health care worker talk with you about any of the things listed below? Please count only discussions, not reading materials or videos. For each item, circle **Y** (Yes) if someone talked with you about it or circle **N** (No) if no one talked with you about it.**

	No	Yes
a. How smoking during pregnancy could affect my baby . . . . .	N	Y
b. Breastfeeding my baby . . . . .	N	Y
c. How drinking alcohol during pregnancy could affect my baby . . . . .	N	Y
d. Using a seat belt during my pregnancy . . . . .	N	Y
e. Birth control methods to use after my pregnancy . . . . .	N	Y
f. Medicines that are safe to take during my pregnancy . . . . .	N	Y
g. How using illegal drugs could affect my baby . . . . .	N	Y
h. Doing tests to screen for birth defects or diseases that run in my family . . . . .	N	Y
i. What to do if my labor starts early . . . . .	N	Y
j. Getting tested for HIV (the virus that causes AIDS) . . . . .	N	Y
k. Physical abuse to women by their husbands or partners . . . . .	N	Y

**22. During any of your prenatal care visits, did a doctor, nurse, or other health care worker ask if you were smoking cigarettes?**

- No
- Yes

**23. During any of your prenatal care visits, did a doctor, nurse, or other health care worker ask if you were drinking alcoholic beverages (beer, wine, wine cooler, or liquor)?**

- No
- Yes

**24. During any of your prenatal care visits, did a doctor, nurse, or other health care worker talk with you about how eating fish containing high levels of mercury could affect your baby?**

- No  
 Yes

**25. At any time during your most recent pregnancy or delivery, did you have a test for HIV (the virus that causes AIDS)?**

- No  
 Yes  
 I don't know

**The next questions are about your most recent pregnancy and things that might have happened during your pregnancy.**

**26. During your most recent pregnancy, were you on WIC (the Special Supplemental Nutrition Program for Women, Infants, and Children)?**

- No  
 Yes

**27. Did you have any of these problems during your most recent pregnancy?** For each item, circle **Y** (Yes) if you had the problem or circle **N** (No) if you did not.

	No	Yes
a. High blood sugar (diabetes) that started <i>before</i> this pregnancy . . . . .	N	Y
b. High blood sugar (diabetes) that started <i>during</i> this pregnancy . . . . .	N	Y
c. Vaginal bleeding . . . . .	N	Y
d. Kidney or bladder (urinary tract) infection . . . . .	N	Y
e. Severe nausea, vomiting, or dehydration . . . . .	N	Y
f. Cervix had to be sewn shut (incompetent cervix) . . . . .	N	Y
g. High blood pressure, hypertension (including pregnancy-induced hypertension [PIH]), preeclampsia, or toxemia . . . . .	N	Y
h. Problems with the placenta (such as abruptio placentae or placenta previa) . . . . .	N	Y
i. Labor pains more than 3 weeks before my baby was due (preterm or early labor) . . . . .	N	Y
j. Water broke more than 3 weeks before my baby was due (premature rupture of membranes [PROM]) . . . . .	N	Y
k. I had to have a blood transfusion . . . . .	N	Y
l. I was hurt in a car accident . . . . .	N	Y

**If you did not have any of these problems, go to Page 6, Question 29.**

**28. Did you do any of the following things because of these problems?** For each item, circle **Y** (Yes) if you did that thing or circle **N** (No) if you did not.

	<b>No</b>	<b>Yes</b>
a. I went to the hospital or emergency room and stayed less than 1 day . . . . .	N	Y
b. I went to the hospital and stayed 1 to 7 days . . . . .	N	Y
c. I went to the hospital and stayed more than 7 days . . . . .	N	Y
d. I stayed in bed at home more than 2 days because of my doctor's or nurse's advice . . . . .	N	Y

**The next questions are about smoking cigarettes and drinking alcohol.**

**29. Have you smoked at least 100 cigarettes in the past 2 years?** (A pack has 20 cigarettes.)

- No —————> **Go to Question 33**  
 Yes

**30. In the 3 months before you got pregnant, how many cigarettes did you smoke on an average day?** (A pack has 20 cigarettes.)

- 41 cigarettes or more  
 21 to 40 cigarettes  
 11 to 20 cigarettes  
 6 to 10 cigarettes  
 1 to 5 cigarettes  
 Less than 1 cigarette  
 None (0 cigarettes)

**31. In the last 3 months of your pregnancy, how many cigarettes did you smoke on an average day?** (A pack has 20 cigarettes.)

- 41 cigarettes or more  
 21 to 40 cigarettes  
 11 to 20 cigarettes  
 6 to 10 cigarettes  
 1 to 5 cigarettes  
 Less than 1 cigarette  
 None (0 cigarettes)

**32. How many cigarettes do you smoke on an average day now?** (A pack has 20 cigarettes.)

- 41 cigarettes or more  
 21 to 40 cigarettes  
 11 to 20 cigarettes  
 6 to 10 cigarettes  
 1 to 5 cigarettes  
 Less than 1 cigarette  
 None (0 cigarettes)

**33. Have you had any alcoholic drinks in the past 2 years?** (A drink is 1 glass of wine, wine cooler, can or bottle of beer, shot of liquor, or mixed drink.)

- No —————> **Go to Question 36**  
 Yes

**34a. During the 3 months before you got pregnant, how many alcoholic drinks did you have in an average week?**

- 14 drinks or more a week  
 7 to 13 drinks a week  
 4 to 6 drinks a week  
 1 to 3 drinks a week  
 Less than 1 drink a week  
 I didn't drink then



**34b. During the 3 months before you got pregnant, how many times did you drink 5 alcoholic drinks or more in one sitting?**

- 6 or more times  
 4 to 5 times  
 2 to 3 times  
 1 time  
 I didn't have 5 drinks or more in 1 sitting  
 I didn't drink then

**35a. During the last 3 months of your pregnancy, how many alcoholic drinks did you have in an average week?**

- 14 drinks or more a week  
 7 to 13 drinks a week  
 4 to 6 drinks a week  
 1 to 3 drinks a week  
 Less than 1 drink a week  
 I didn't drink then

**35b. During the last 3 months of your pregnancy, how many times did you drink 5 alcoholic drinks or more in one sitting?**

- 6 or more times  
 4 to 5 times  
 2 to 3 times  
 1 time  
 I didn't have 5 drinks or more in 1 sitting  
 I didn't drink then

**Pregnancy can be a difficult time for some women. These next questions are about things that may have happened before and during your most recent pregnancy.**

**36. This question is about things that may have happened during the 12 months before your new baby was born.** For each item, circle **Y** (Yes) if it happened to you or circle **N** (No) if it did not. (It may help to use the calendar.)

	No	Yes
a. A close family member was very sick and had to go into the hospital. . . .	N	Y
b. I got separated or divorced from my husband or partner . . . . .	N	Y
c. I moved to a new address . . . . .	N	Y
d. I was homeless . . . . .	N	Y
e. My husband or partner lost his job . . .	N	Y
f. I lost my job even though I wanted to go on working. . . . .	N	Y
g. I argued with my husband or partner more than usual. . . . .	N	Y
h. My husband or partner said he didn't want me to be pregnant. . . . .	N	Y
i. I had a lot of bills I couldn't pay. . . .	N	Y
j. I was in a physical fight . . . . .	N	Y
k. My husband or partner or I went to jail . . . . .	N	Y
l. Someone very close to me had a bad problem with drinking or drugs . . . .	N	Y
m. Someone very close to me died . . . .	N	Y

**37. During the 12 months before your new baby was born, did you ever eat less than you felt you should because there wasn't enough money to buy food?**

- No  
 Yes

The next questions are about the time during the *12 months before* you got pregnant with your new baby.

**38a.** During the *12 months before* you got pregnant, did an ex-husband or ex-partner push, hit, slap, kick, choke, or physically hurt you in any other way?

- No  
 Yes

**38b.** During the *12 months before* you got pregnant, were you physically hurt in any way by your husband or partner?

- No  
 Yes

The next questions are about the time during your most recent pregnancy.

**39a.** During your most recent pregnancy, did an ex-husband or ex-partner push, hit, slap, kick, choke, or physically hurt you in any other way?

- No  
 Yes

**39b.** During your most recent pregnancy, were you physically hurt in any way by your husband or partner?

- No  
 Yes

The next questions are about your labor and delivery. (It may help to look at the calendar when you answer these questions.)

**40.** When was your baby due?

Month Day Year

**41.** When did you go into the hospital to have your baby?

Month Day Year

I didn't have my baby in a hospital

**42.** When was your baby born?

Month Day Year

**43.** When were you discharged from the hospital after your baby was born? (It may help to use the calendar.)

Month Day Year

I didn't have my baby in a hospital

**44. How was your delivery paid for?****Check all that apply**

- Oregon Health Plan or Medicaid
- Personal income (cash, check, or credit card)
- Health insurance or HMO (including insurance from your work or your husband's work)
- Indian Health Service
- Other —————> Please tell us:

---

**The next questions are about the time since your new baby was born.**

**45. After your baby was born, was he or she put in an intensive care unit?**

- No
- Yes
- I don't know

**46. After your baby was born, how long did he or she stay in the hospital?**

- Less than 24 hours (less than 1 day)
- 24 to 48 hours (1 to 2 days)
- 3 days
- 4 days
- 5 days
- 6 days or more
- My baby was not born in a hospital
- My baby is still in the hospital —————>

**Go to Question 49****47. Is your baby alive now?**

- No —————> **Go to Page 11, Question 60**
- Yes

**48. Is your baby living with you now?**

- No —————> **Go to Page 11, Question 60**
- Yes

**49. Did you ever breastfeed or pump breast milk to feed your new baby after delivery?**

- No —————> **Go to Page 10, Question 54**
- Yes

**50. Are you still breastfeeding or feeding pumped milk to your new baby?**

- No
- Yes —————> **Go to Page 10, Question 53**

**51. How many weeks or months did you breastfeed or pump milk to feed your baby?**

Weeks **OR**  Months

- Less than 1 week

**52. What were your reasons for stopping breastfeeding?**

Check all that apply

- My baby had difficulty nursing
- Breast milk alone did not satisfy my baby
- I thought my baby was not gaining enough weight
- My baby got sick and could not breastfeed
- My nipples were sore, cracked, or bleeding
- I thought I was not producing enough milk
- I had too many other household duties
- I felt it was the right time to stop breastfeeding
- I got sick and could not breastfeed
- I went back to work or school
- I wanted or needed someone else to feed the baby
- My baby was jaundiced (yellowing of the skin or whites of the eyes)
- Other —————> Please tell us:

---

**53. How old was your baby the first time you fed him or her anything besides breast milk? Include formula, baby food, juice, cow's milk, water, sugar water, or anything else you fed your baby.**

\_\_\_\_\_ Weeks **OR** \_\_\_\_\_ Months

- My baby was less than 1 week old
- I have not fed my baby anything besides breast milk

**If your baby was not born in a hospital, go to Question 55.**

**54. This question asks about things that may have happened at the hospital where your new baby was born. For each item, circle Y (Yes) if it happened or circle N (No) if it did not happen.**

	No	Yes
a. Hospital staff gave me information about breastfeeding . . . . .	N	Y
b. My baby stayed in the same room with me at the hospital . . . . .	N	Y
c. I breastfed my baby in the hospital . . . . .	N	Y
d. I breastfed my baby in the first hour after my baby was born. . . . .	N	Y
e. Hospital staff helped me learn how to breastfeed . . . . .	N	Y
f. My baby was fed only breast milk at the hospital . . . . .	N	Y
g. Hospital staff told me to breastfeed whenever my baby wanted . . . . .	N	Y
h. The hospital gave me a gift pack with formula . . . . .	N	Y
i. The hospital gave me a telephone number to call for help with breastfeeding . . . . .	N	Y
j. My baby used a pacifier in the hospital . . . . .	N	Y

**If your baby is still in the hospital, go to Question 60.**

**55. About how many hours a day, on average, is your new baby in the same room with someone who is smoking?**

\_\_\_\_\_ Hours

- Less than 1 hour a day
- My baby is never in the same room with someone who is smoking

**56. How do you *most often* lay your baby down to sleep now?**

Check one answer

- On his or her side  
 On his or her back  
 On his or her stomach

**57. How often does your new baby sleep in the same bed with you or anyone else?**

- Always  
 Often  
 Sometimes  
 Rarely  
 Never

**58. Was your new baby seen by a doctor, nurse, or other health care worker during the first week after he or she left the hospital?**

- No  
 Yes

**59. Has your new baby had a well-baby checkup?**  
 (A well-baby checkup is a regular health visit for your baby usually at 2, 4, or 6 months of age.)

- No  
 Yes

**60. Are you or your husband or partner doing anything *now* to keep from getting pregnant?**  
 (Some things people do to keep from getting pregnant include not having sex at certain times [rhythm] or withdrawal, and using birth control methods such as the pill, condoms, cervical ring, IUD, having their tubes tied, or their partner having a vasectomy.)

- No  
 Yes →

Go to Question 62

**61. What are your or your husband's or partner's reasons for not doing anything to keep from getting pregnant *now*?**

Check all that apply

- I am not having sex  
 I want to get pregnant  
 I don't want to use birth control  
 My husband or partner doesn't want to use anything  
 I don't think I can get pregnant (sterile)  
 I can't pay for birth control  
 I am pregnant now  
 Other → Please tell us:

If you or your husband or partner is not doing anything to keep from getting pregnant *now*, go to Page 12, Question 63.

**62. What kind of birth control are you or your husband or partner using *now* to keep from getting pregnant?**

Check all that apply

- Tubes tied or closed (female sterilization)  
 Vasectomy (male sterilization)  
 Pill  
 Condoms  
 Shot once a month (Lunelle®)  
 Shot once every 3 months (Depo-Provera®)  
 Contraceptive patch (OrthoEvra®)  
 Diaphragm, cervical cap, or sponge  
 Cervical ring (NuvaRing® or others)  
 IUD (including Mirena®)  
 Rhythm method or natural family planning  
 Withdrawal (pulling out)  
 Not having sex (abstinence)  
 Other → Please tell us:

The next few questions are about the time during the *12 months before your new baby was born*.

**63. During the 12 months before your new baby was born, what were the sources of your household's income?**

Check all that apply

- Paycheck or money from a job
- Money from family or friends
- Money from a business, fees, dividends, or rental income
- Aid such as Temporary Assistance for Needy Families (TANF), welfare, WIC, public assistance, general assistance, food stamps, or Supplemental Security Income
- Unemployment benefits
- Child support or alimony
- Social security, workers' compensation, disability, veteran benefits, or pensions
- Other ————— ➤ Please tell us:

**64. During the 12 months before your new baby was born, what was your total household income before taxes?** Include your income, your husband's or partner's income, and any other income you may have used. (All information will be kept private and will not affect any services you are now getting.)

Check one answer

- Less than \$10,000
- \$10,000 to \$14,999
- \$15,000 to \$19,999
- \$20,000 to \$24,999
- \$25,000 to \$34,999
- \$35,000 to \$49,999
- \$50,000 or more

**65. During the 12 months before your new baby was born, how many people, including yourself, depended on this income?**

People

The remaining questions are on a variety of topics of importance to programs for Oregon mothers and babies. Remember that your answers should be about your *most recent pregnancy with your new baby*.

**66. Before you got pregnant with your new baby, had you ever heard or read about emergency birth control (the "morning-after pill")?** This combination of pills is used to prevent pregnancy up to 3 days after unprotected sex.

- No
- Yes

If you or your husband or partner was using birth control when you got pregnant with your new baby, go to Question 68a.

**67. When you got pregnant with your new baby, would you have used a birth control method if you had insurance that paid for it?**

- No
- Yes

**68a. While you were pregnant, how often did you feel down, depressed, or hopeless?**

- Always
- Often
- Sometimes
- Rarely
- Never

**68b. While you were pregnant, how often did you have little interest or little pleasure in doing things?**

- Always  
 Often  
 Sometimes  
 Rarely  
 Never

**69. During your most recent pregnancy, did you receive any of the following services?** For each one, circle **Y** (Yes) if you received the service or circle **N** (No) if you did not receive the service.

**Did you receive—**

	No	Yes
a. Help with an alcohol or drug problem. . . . .	N	Y
b. Help to reduce violence in your home . . . . .	N	Y
c. Counseling information for family and personal problems . . . . .	N	Y
d. Help to quit smoking. . . . .	N	Y

**70. During any of your prenatal care visits or after your most recent delivery, did a doctor, nurse, or other health care worker ever advise you to quit smoking?**

- Yes, during my prenatal care visits  
 Yes, after my delivery  
 Yes, both times  
 No  
 No, I did not smoke at that time

**If your baby is no longer alive or is not living with you, go to Question 74.**

**71. During any of your prenatal care visits or after your most recent delivery, did a doctor, nurse, or other health care worker talk with you about how secondhand smoke could affect your baby after birth?**

- Yes, during my prenatal care visits  
 Yes, after my delivery  
 Yes, both times  
 No

**72. After your new baby was born, did a doctor, nurse, or other health care worker talk with you about how to prevent your baby from getting tooth decay?**

- No  
 Yes

**73a. Do you ever put your baby to bed with a bottle?**

- No —————> **Go to Question 74**  
 Yes

**73b. What do you put in the bottles that your baby takes to bed?**

**Check all that apply**

- Water  
 Something other than water

**74. In the past month, how many days a week did you get at least 30 minutes of physical activity or exercise?** (For example, walking, dancing, yard work, or sweeping.)

- Less than 1 day per week  
 1 to 4 days per week  
 5 or more days per week

**75a. Since your new baby was born, how often have you felt down, depressed, or hopeless?**

- Always
- Often
- Sometimes
- Rarely
- Never

**75b. Since your new baby was born, how often have you had little interest or little pleasure in doing things?**

- Always
- Often
- Sometimes
- Rarely
- Never

**76. This question is about the care of your teeth during your most recent pregnancy.** For each item, circle **Y** (Yes) if it is true or circle **N** (No) if it is not true.

**No    Yes**

- a. I needed to see a dentist for a problem . . . . . N    Y
- b. I went to a dentist or dental clinic. . . . N    Y
- c. A dental or other health care worker talked with me about how to care for my teeth and gums . . . . . N    Y

**77. How long has it been since you had your teeth cleaned by a dentist or a dental hygienist?**

- Within the past year (less than 12 months)
- 1 to less than 2 years (12 to 23 months)
- 2 to less than 5 years (24 to 59 months)
- 5 or more years (60 or more months)
- Never

**78. Not including yourself, is there anyone in your household who smokes cigarettes, cigars, or pipes?**

- No
- Yes

**79. Which of the following statements best describes the rules about smoking *inside* your home now?**

**Check one answer**

- No one is allowed to smoke anywhere inside my home
- Smoking is allowed in some rooms or at some times
- Smoking is permitted anywhere inside my home

**80. What is today's date?**

<input style="width: 100%; height: 20px;" type="text"/>	<input style="width: 100%; height: 20px;" type="text"/>	<input style="width: 100%; height: 20px;" type="text"/>
Month	Day	Year



**Please use this space for any additional comments you would like to make  
about the health of mothers and babies in Oregon.**

*Thanks for answering our questions!*

*Your answers will help us work to make Oregon  
mothers and babies healthier.*

## **Appendix B**

2006 Oregon PRAMS-2

(longitudinal follow-up survey of 2004 PRAMS)

*In this first part of the survey, we would like to ask some questions about YOU. Please check the box next to your answer, fill in the blank, or circle as directed.*

**1. What is your date of birth?**

\_\_\_\_\_ 19\_\_\_\_\_  
 Month Day Year

**2. What is the highest level of school you have completed?**  
*Check one answer.*

- Less than 12<sup>th</sup> grade
- 12<sup>th</sup> grade or GED
- More than 12<sup>th</sup> grade

**3a. What is your current marital status?**  
*Check one answer.*

- Never married
- Married
- Widowed
- Divorced
- Separated

**3b. Are you living with .....?**  
*Check all that apply.*

- Your spouse or partner
- Other adult (not spouse or partner)
- No other adult(s)

**4a. Have you lived in the United States all your life?**

- No
- Yes → Go to Question 5

**4b. For how many years have you lived in the United States?**

- 0 to 3 years
- 4 to 6 years
- 7 to 13 years
- 14 to 20 years
- More than 20 years

**5. Are you employed?**  
*Check one answer.*

- Yes, full time
- Yes, part time
- No, but I am looking for work
- No, I am not looking for work

**6. Is your spouse or partner, who is living with you, employed?**  
*Check one answer.*

- Yes, full time
- Yes, part time
- No, but they are looking for work
- No, they are not looking for work
- I do not have a spouse or partner living with me

*The next questions are about your health insurance and medical history.*

**7. What kind of health insurance do you have right now?**  
*Check all that apply.*

- I don't have insurance
- Oregon Health Plan (OHP), Medicaid or SCHIP
- Medicare
- Private Insurance
- Military/CHAMPUS
- Indian Health Service
- Other → Please tell us: \_\_\_\_\_

**8. During any of your health care visits in the last 12 months, did a doctor, nurse, or other health care worker talk with you about any of the things listed below?** Please count only discussions, not reading materials or videos. For each item, circle **Y** (Yes) if someone talked with you about it, circle **N** (No) if no one talked with you about it or if it did not apply to you.

**Did they....?** **No** **Yes**

- |                                                         |   |   |
|---------------------------------------------------------|---|---|
| a. Talk about physical abuse to women by their partners | N | Y |
| b. Ask you if you smoked                                | N | Y |
| c. Advise you to quit smoking                           | N | Y |
| d. Offer you help on how to quit smoking                | N | Y |
| e. Talk about how drinking alcohol can affect you       | N | Y |

**9. Have you ever been told by a doctor, nurse or other health care worker that you had diabetes (sugar diabetes) during any of your pregnancies?**

- No
- Yes

**10. Have you ever been told by a doctor, nurse or other health care worker that you had diabetes (sugar diabetes) when you were not pregnant?**

- No
- Yes

**11. Have any of your family members ever been told by a doctor, nurse or other health care worker that they had diabetes (sugar diabetes)?**

For each family member, circle **Y** (Yes) if they were told that they had diabetes, circle **N** (No) if they were not told, or circle **DK** (Don't Know) if you do not know.

- |                                                                                  | No | Yes | Don't Know |
|----------------------------------------------------------------------------------|----|-----|------------|
| a. Your two-year-old                                                             | N  | Y   | DK         |
| b. Your two-year-old's father                                                    | N  | Y   | DK         |
| c. Your two-year-old's brothers or sisters (including half brothers and sisters) | N  | Y   | DK         |
| d. Your mother                                                                   | N  | Y   | DK         |
| e. Your father                                                                   | N  | Y   | DK         |
| f. Your brothers or sisters                                                      | N  | Y   | DK         |

**12. Have you ever been told by a doctor, nurse or other health care worker that you had asthma?**

- No  
 Yes

**13. Have any of your family members ever been told by a doctor, nurse or other health care worker that they had asthma?** For each family member, circle **Y** (Yes) if they were told that they had asthma, circle **N** (No) if they were not told, or circle **DK** (Don't Know) if you do not know.

- |                                                                                  | No | Yes | Don't Know |
|----------------------------------------------------------------------------------|----|-----|------------|
| a. Your two-year-old                                                             | N  | Y   | DK         |
| b. Your two-year-old's father                                                    | N  | Y   | DK         |
| c. Your two-year-old's brothers or sisters (including half brothers and sisters) | N  | Y   | DK         |
| d. Your mother                                                                   | N  | Y   | DK         |
| e. Your father                                                                   | N  | Y   | DK         |
| f. Your brothers or sisters                                                      | N  | Y   | DK         |

*The next questions are about smoking cigarettes and drinking alcohol.*

**14. Have you smoked at least 100 cigarettes in your entire life?** (A pack has 20 cigarettes.)

- No → Go to Question 16  
 Yes

**15. How many cigarettes do you smoke on an average day now?** (A pack has 20 cigarettes.)

- 41 cigarettes or more  
 21 to 40 cigarettes  
 11 to 20 cigarettes  
 6 to 10 cigarettes  
 1 to 5 cigarettes  
 Less than 1 cigarette  
 None (0 cigarettes)

**16. Not including yourself, is there anyone in your household who smokes cigarettes, cigars, or pipes?**

- No  
 Yes

**17. Which of the following statements best describes the rules about smoking *inside* your home *now*?**

- No one is allowed to smoke anywhere inside my home  
 Smoking is allowed in some rooms or at some times  
 Smoking is permitted anywhere inside my home

**18. In the past 12 months, how many alcoholic drinks did you have in an average week?** (A drink is 1 glass of wine, wine cooler, can or bottle of beer, shot of liquor, or mixed drink.)

- 14 drinks or more a week  
 7 to 13 drinks a week  
 4 to 6 drinks a week  
 1 to 3 drinks a week  
 Less than 1 drink a week  
 I didn't drink then

**19. In the past 12 months, how many times did you drink 4 alcoholic drinks or more in one sitting?**

- 6 or more times  
 4 to 5 times  
 2 to 3 times  
 1 time  
 I didn't have 4 drinks or more in 1 sitting  
 I didn't drink then

**20. Since your two-year-old was born, have you drunk more alcohol than you intended?**

- No  
 Yes

**21. In the past 12 months, have you ever felt the need to cut down on drinking alcohol?**

- No  
 Yes

*The next questions are about emotions and stress.*

**22a. During the FIRST 12 months of your two-year-old's life, was there a period of two or more weeks when almost every day you:**

- |                                                                                | No | Yes |
|--------------------------------------------------------------------------------|----|-----|
| a. Felt sad, blue or depressed for most of the day                             | N  | Y   |
| b. Lost interest or pleasure in most things you usually cared about or enjoyed | N  | Y   |

**22b. In the PAST 12 months, has there been period of two or more weeks when almost every day you:**

- |                                                                                | No | Yes |
|--------------------------------------------------------------------------------|----|-----|
| a. Felt sad, blue or depressed for most of the day                             | N  | Y   |
| b. Lost interest or pleasure in most things you usually cared about or enjoyed | N  | Y   |

**23. In the past 12 months, has a doctor, nurse, or other health care or mental health worker told you that you had:**

- |                                      | No | Yes |
|--------------------------------------|----|-----|
| a. Depression                        | N  | Y   |
| b. Any other mental health condition | N  | Y   |

**24. In the past 12 months, have you taken prescription medications for:**

- |                                      | No | Yes |
|--------------------------------------|----|-----|
| a. Depression                        | N  | Y   |
| b. Any other mental health condition | N  | Y   |

**25. This question is about things that may have happened to you in the past 12 months.** For each item, circle Y (Yes) if it happened to you or circle N (No) if it did not.

- |                                                                        | No | Yes |
|------------------------------------------------------------------------|----|-----|
| a. A close family member was very sick and had to go into the hospital | N  | Y   |
| b. I was very sick                                                     | N  | Y   |
| c. I got separated or divorced from my spouse or partner               | N  | Y   |
| d. I moved to a new address                                            | N  | Y   |
| e. I was homeless                                                      | N  | Y   |
| f. My spouse or partner lost his or her job                            | N  | Y   |
| g. I lost my job even though I wanted to go on working                 | N  | Y   |
| h. I argued with my spouse or partner more than usual                  | N  | Y   |
| i. I had a lot of bills I couldn't pay                                 | N  | Y   |
| j. I was in a physical fight                                           | N  | Y   |
| k. My spouse or partner or I went to jail                              | N  | Y   |
| l. Someone very close to me had a bad problem with drinking or drugs   | N  | Y   |
| m. Someone very close to me died                                       | N  | Y   |

**26. In the past 12 months, did an intimate partner (current or former spouse, boyfriend, girlfriend, or date) do any of the following to you?** For each item, circle Y (Yes) if it happened to you or circle N (No) if it did not.

- |                                                                                                 | No | Yes |
|-------------------------------------------------------------------------------------------------|----|-----|
| a. Yelled and screamed at you, threatened you or made you feel unsafe                           | N  | Y   |
| b. Tried to limit your contact with family or friends                                           | N  | Y   |
| c. Prevented you from knowing about or having access to your shared income, even when you asked | N  | Y   |
| d. Pushed, hit, slapped, kicked, choked, or physically hurt you in any other way                | N  | Y   |
| e. Had sex with you against your will or without your consent                                   | N  | Y   |

**27. For each of the following items, circle Y (Yes) if it describes your current situation or circle N (No) if it does not.**

- |                                                                                              | No | Yes |
|----------------------------------------------------------------------------------------------|----|-----|
| a. You have someone who would loan you money for food or bills if you needed it              | N  | Y   |
| b. You have someone who would help you if you were sick and needed to be in bed              | N  | Y   |
| c. You have someone who would take you to the clinic or doctor's office if you needed a ride | N  | Y   |
| d. You have someone you can count on to listen to you when you need to talk                  | N  | Y   |
| e. You have someone who shows you love and affection                                         | N  | Y   |

*The next questions are about pregnancy and birth control.*

**28. Have you been pregnant since your two-year-old was born?** (If you are currently pregnant, count this pregnancy too.)

- No → Go to Question 30  
 Yes

**29. Thinking back to just before your most recent pregnancy, how did you feel about becoming pregnant?**  
*Check one answer.*

- I wanted to be pregnant sooner  
 I wanted to be pregnant later  
 I wanted to be pregnant then  
 I didn't want to be pregnant then or at any time in the future

**30. Are you or your spouse or partner doing anything now to keep from getting pregnant?** (Some things people do to keep from getting pregnant include having their tubes tied or their partner having a vasectomy, not having sex at certain times [rhythm] or withdrawal, and using

birth control methods such as the pill, condoms, the patch, shots, or IUDs.)

- No
- Yes → **Go to Question 32**

**31. What are you or your spouse's or partner's reasons for not doing anything to keep from getting pregnant now?**  
*Check all that apply.*

- I am not having sex
- I want to get pregnant
- I am pregnant now
- I am breastfeeding
- I don't want to use birth control
- My spouse or partner doesn't want to use birth control
- I don't think I can get pregnant (sterile)
- I can't pay for birth control
- Same-sex partner
- Other → Please tell us:

**If you or your spouse/partner are not doing anything to keep from getting pregnant now, go to Question 33.**

**32. What kind of birth control are you or your spouse/partner using now to keep from getting pregnant?**  
*Check all that apply.*

- Tubes tied or closed (female sterilization)
- Vasectomy (male sterilization)
- Pill
- Condoms
- Shot once every 3 months (Depo-Provera®)
- Contraceptive patch (OrthoEvra®)
- Diaphragm, cervical cap, or sponge
- Vaginal ring (NuvaRing®)
- IUD (including Mirena®)
- Rhythm method or natural family planning
- Withdrawal (pulling out)
- Not having sex (abstinence)
- Other → Please tell us:

***The next questions are about your current activities.***

**33. How many times per week do you take a multivitamin?** These are pills that contain many different vitamins and minerals.

- I don't take a multivitamin at all
- 1 to 3 times a week
- 4 to 6 times a week
- Every day of the week

**34. In the past month, how many days a week did you get at least 30 minutes of physical activity or exercise?** (For example, walking, dancing, yard work, or sweeping.)

- Less than 1 day per week
- 1 to 4 days per week
- 5 or more days per week

**35. In the past 12 months, did you ever eat less than you felt you should because there wasn't enough money to buy food?**

- No
- Yes

**36. Is the tap water in your home fluoridated?**

- No
- Yes
- I don't know

**37. How much do you weigh now?**

\_\_\_\_\_ Pounds **OR** \_\_\_\_\_ Kilos

**38. What do you think about your weight?**  
*Check one answer.*

- I am underweight
- I am about the right weight
- I am overweight

***The next questions are general questions.***

**39. In the past 12 months, have you or your two-year-old needed or received any of the following?**

	Didn't need it	Needed it, didn't get it	Needed it, got it
a. WIC Services	DN	N	NG
b. Food Stamps or money to buy food	DN	N	NG
c. Other financial assistance (for example, AFDC, TANF, subsidized rent, etc.)	DN	N	NG
d. Help with an alcohol or drug problem	DN	N	NG
e. Help to stop smoking	DN	N	NG
f. Help with transportation	DN	N	NG
g. Help paying for education or job training	DN	N	NG
h. Help with a family violence problem	DN	N	NG
i. Help or counseling for other family or personal problems	DN	N	NG

**40. What is your total annual household income before taxes?** Include your income, your spouse's/partner's income, and any other income you may have. *(All information will be kept private and will not affect any services you are now getting.)*  
**Check one answer.**

- Less than \$10,000
- \$10,000 to \$14,999
- \$15,000 to \$19,999
- \$20,000 to \$24,999
- \$25,000 to \$29,999
- \$30,000 to \$34,999
- \$35,000 to \$49,999
- \$50,000 or more

**41. How many people, including yourself, depend on this income?**

\_\_\_\_\_ People

*In this last part of the survey are questions about your two-year-old-child.*

**42. What is your two-year-old's date of birth?**

\_\_\_\_\_ 20\_\_\_\_\_  
Month Day Year

**43. Is your two-year-old alive now?**

- Yes → **Go to Question 44**
- No

*If your child is no longer alive, we are truly sorry about your loss and extend our sympathy to you and your family. The answers you have given are especially important and could help us learn about ways to improve the health and safety of children in the future.*

**When did your child die?**

\_\_\_\_\_ 20\_\_\_\_\_  
Month Day Year

**If your child is no longer alive, thank you for answering these questions. Please provide today's date on page 14, Question 92.**

**44. Is your two-year-old living with you now?**

- No
- Yes → **Go to Question 45a**

**If your two-year-old is not living with you, thank you for answering these questions. Please provide today's date on page 14, Question 92.**

**45a. How much does your two-year-old weigh?**

\_\_\_\_\_ Pounds **OR** \_\_\_\_\_ Kilos

**45b. How do you know your child's weight?**  
**Check one answer.**

- Measured by health care provider  
(Approximate Date: \_\_\_\_\_)
- Measured by someone else  
(Approximate Date: \_\_\_\_\_)
- Estimated now
- Other → Please tell us:  
\_\_\_\_\_

**46a. How tall is your two-year-old?**

\_\_\_\_\_ Feet \_\_\_\_\_ Inches

**OR** \_\_\_\_\_ Centimeters

**46b. How do you know your child's height?**  
**Check one answer.**

- Measured by health care provider  
(Approximate Date: \_\_\_\_\_)
- Measured by someone else  
(Approximate Date: \_\_\_\_\_)
- Estimated now
- Other → Please tell us:  
\_\_\_\_\_

**47. How would you rate your two-year-old's health in general?**

**Check one answer.**

- Excellent
- Very Good
- Good
- Fair
- Poor

*The next questions are about breastfeeding.*

**48. Did you ever breastfeed or pump breast milk to feed your child, who is now two-year's-old?**

- No → **Go to Page 8, Question 52**
- Yes

**49. During the first 12 months, which of the following helped you to continue breastfeeding your two-year-old for as long as you did?**  
**Check all that apply.**

- Support from friends and family
  - Support from my employer
  - Support from a health care provider
  - Convenience to me
  - Cost savings
  - Benefits to my child
  - Benefits for myself
  - My own commitment to breastfeed
  - My baby was not ready to stop breastfeeding
  - Other → Please tell us:
- 

**50. How old was your two-year-old when he/she completely stopped breastfeeding?**

\_\_\_\_\_ months old

- Still breastfeeding → **Go to Page 8, Question 52**

**51. What were your reasons for stopping breastfeeding?**

*Check all that apply.*

- I felt it was the right time to stop breastfeeding
  - I went back to work or school
  - There was no place to pump or feed my child at work/school
  - My child weaned himself/herself
  - My child became sick and could not breastfeed
  - I wanted or needed someone else to feed my child
  - My child's teeth came in
  - My child seemed too old to breastfeed
  - I became sick and could not breastfeed
  - I thought my child was not gaining enough weight
  - I thought I wasn't producing enough milk
  - I had too many other responsibilities
  - Family or friends suggested that I stop breastfeeding
  - My doctor suggested that I stop breastfeeding
  - I believed that my milk became less nutritious as my child got older
  - Other → Please tell us:
- 

*The next questions are about your two-year-old's eating habits now.*

**52. What do you think about the amount your two-year-old eats?**

*Check one answer.*

- My child does not eat enough
- My child eats the right amount
- My child eats too much

**53. How many days in a typical week does your two-year-old eat each of the foods listed below?**

*Circle the number of days.*

Vegetables other than potatoes	0 1 2 3 4 5 6 7 days
French Fries	0 1 2 3 4 5 6 7 days
Fresh or canned fruit	0 1 2 3 4 5 6 7 days
Candy or cookies	0 1 2 3 4 5 6 7 days

**54. How many days in a typical week does your two-year-old drink the following drinks?**

*Circle the number of days.*

Milk	0 1 2 3 4 5 6 7 days
Fruit juices	0 1 2 3 4 5 6 7 days
Fruit drinks & Kool-Aid	0 1 2 3 4 5 6 7 days
Soda pop	0 1 2 3 4 5 6 7 days
Plain water	0 1 2 3 4 5 6 7 days
Sports drinks (example: Gatorade, PowerAde)	0 1 2 3 4 5 6 7 days

**55. In the past week, how many days did your two-year-old eat restaurant, fast food or take-out food?**

Take-out food could be from a restaurant, supermarket or deli counter.

*Circle the number of days.*

0 1 2 3 4 5 6 7 days

**56. Have you changed the amount or type of fish your child eats, due to advice you have read, seen or heard about mercury in fish?**

- No
- Yes
- I am not aware of this advice

**57a. Do you currently ever put your two-year-old to bed with a bottle?**

- No → **Go to Question 58**
- Yes

**57b. What do you put in the bottles that your two-year-old takes to bed?**

*Check all that apply.*

- Water
- Something other than water

**58. Does your family eat meals together?**

*Check one answer.*

- Always
- Usually
- Sometimes
- Never

**59. Has your two-year-old ever been on WIC (the Special Supplemental Nutrition Program for Women, Infants, and Children)?**



- No
- Yes, on WIC now
- Yes, but no longer on WIC

**60. What has a doctor, nurse or other health care worker told you about your two-year-old's weight?**

*Check one answer.*

- That s/he is underweight
- That their weight is normal
- That s/he is overweight, but that you shouldn't worry about it
- That s/he is overweight, and that it is a problem
- Other → Please tell us: \_\_\_\_\_
- They have not talked to me about my child's weight

**61. What do you think about your two-year-old's weight?**

*Check one answer.*

- My child is underweight
- My child is about the right weight
- My child is overweight

*The next questions are about your two-year-old's health insurance and health care.*

**62. What kind of health insurance did your two-year-old have 12 months ago (at 1 year of age)?**

*Check all that apply.*

- None
- Oregon Health Plan (OHP), Medicaid or SCHIP
- Medicare
- Private Insurance
- Military/CHAMPUS
- Indian Health Service
- Other → Please tell us: \_\_\_\_\_

**63. What kind of health insurance does your two-year-old have now?**

*Check all that apply.*

- None
- Oregon Health Plan (OHP), Medicaid or SCHIP
- Medicare
- Private Insurance
- Military/CHAMPUS
- Indian Health Service
- Other → Please tell us: \_\_\_\_\_

**64. Since he or she was born, has there ever been a time when your two-year-old did not have medical insurance?**

- No
- Yes

**65. Does your two-year-old have a regular health care provider now?**

- No
- Yes

**66. Since your two-year-old was 12 months old, how many of his/her doctor or health care provider visits were for well-child care or immunizations? (Well-child care visits are not for sickness or injuries.)**

\_\_\_\_\_ Visits

- My two-year-old has not had any well-child or immunization visits. → **Go to Page 10, Question 68**

**67. What kind of health care provider does your two-year-old see most of the time for well-child care visits?**

*Check one answer.*

- Family doctor (family practice or general practitioner)
- Pediatrician
- Physician's assistant
- Nurse practitioner (PNP, FNP)
- Naturopath, Homeopath
- Other → Please tell us: \_\_\_\_\_

**68. Here is a list of problems some people have getting health care for their children.** For each item, circle **Y** (Yes) if it was a problem for you or circle **N** (No) if it was not a problem or did not apply to you.

	<b>No</b>	<b>Yes</b>
a. I couldn't get an appointment when I wanted one	N	Y
b. I didn't have enough money or insurance to pay for the visits	N	Y
c. I had no way to get to the clinic or doctor's office	N	Y
d. I couldn't take time off from work	N	Y
e. My child didn't have a regular health care provider to go to	N	Y
f. I couldn't find a provider who would take my child	N	Y
g. The services my child needed weren't available in my community	N	Y
h. I had no one to take care of my other children	N	Y
i. My child's health care provider didn't		

- think s/he needed services N Y  
 j. I had too many other things going on N Y  
 k. Other → Please tell us: N Y

**69. During any of your two-year-old's health care visits, did a doctor, dentist, nurse, or other health care worker talk with you about any of the things listed below?** Please count only discussions, not reading materials or videos. For each item, circle **Y** (Yes) if someone talked with you about it or circle **N** (No) if no one talked with you about it.

- |                                                                            | No | Yes |
|----------------------------------------------------------------------------|----|-----|
| a. Your child's nutrition and feeding                                      | N  | Y   |
| b. Using a car seat                                                        | N  | Y   |
| c. Your child's teeth and dental health                                    | N  | Y   |
| d. How your child is growing and developing                                | N  | Y   |
| e. Your child's vision and hearing                                         | N  | Y   |
| f. Things you can do to help your child learn and grow                     | N  | Y   |
| a. Your child's social and emotional health                                | N  | Y   |
| h. Your child's behavior                                                   | N  | Y   |
| i. Physical activity and exercise for your child                           | N  | Y   |
| j. Places you could take your child for other services                     | N  | Y   |
| k. Questions or concerns you have about your child                         | N  | Y   |
| l. Immunizations (baby shots)                                              | N  | Y   |
| m. Sleeping and naptime behaviors                                          | N  | Y   |
| n. How secondhand smoke could affect your child's health                   | N  | Y   |
| o. How eating fish containing high levels of mercury can affect your child | N  | Y   |
| p. Preventing lead poisoning                                               | N  | Y   |
| q. Your child's weight                                                     | N  | Y   |
| r. How to care for your two year-old's teeth and gums                      | N  | Y   |
| s. The use of fluoride drops or tablets in your home                       | N  | Y   |
| t. Fluoride varnish application                                            | N  | Y   |
| u. Assisting your child in brushing his/her teeth                          | N  | Y   |
| v. Fluoride in your tap water                                              | N  | Y   |

*The next questions are about your two-year-old's immunizations or shots against childhood diseases.*

**70. Has your two-year-old ever been given any immunizations or baby shots?**  
*Check one answer.*

- Yes, all recommended shots  
 Yes, some recommended shots  
 No, none

**71a. Have you ever received a reminder for your two-year-old's immunization shots?** A reminder could include postcards, letters or phone calls.

- No → Go to Question 72  
 Yes  
 I don't know → Go to Question 72

**71b. From whom did you receive the reminder?**  
*Check all that apply.*

- Doctor's office  
 HMO, health plan or insurance  
 County health department  
 Oregon Immunization ALERT  
 Other → Please tell us:  


---

 I don't remember

**72. Here is a list of reasons people can have to delay or prevent them from getting their child's shots or immunizations.** For each item, circle **Y** (Yes) if it was ever a reason you didn't get your two-year-old's shots or circle **N** (No) if it was not a reason or did not apply to you.

- |                                                                                | No | Yes |
|--------------------------------------------------------------------------------|----|-----|
| a. I didn't have childcare for other children                                  | N  | Y   |
| b. I couldn't get an appointment                                               | N  | Y   |
| c. I couldn't find doctor or clinic hours when I was able to go                | N  | Y   |
| d. I was referred to other health care providers or clinics for shots          | N  | Y   |
| e. I couldn't afford a health care visit                                       | N  | Y   |
| f. I couldn't afford the cost of shots                                         | N  | Y   |
| g. I wanted to wait until my child was older for some shots                    | N  | Y   |
| h. My child's health care provider told us to wait on some shots that were due | N  | Y   |
| i. I thought my child was too sick to get shots                                | N  | Y   |
| j. I didn't have transportation                                                | N  | Y   |
| k. I didn't know <i>when</i> the shots were due                                | N  | Y   |
| l. I didn't know <i>where</i> to go for shots                                  | N  | Y   |
| m. I couldn't take time off from work or school                                | N  | Y   |
| n. I didn't think about getting the shots                                      | N  | Y   |
| o. I didn't get around to getting the shots                                    | N  | Y   |
| p. Other → Please tell us:                                                     | N  | Y   |

**73. Here is a list of concerns people may have with immunizations or shots recommended for their two-year-olds.** For each item, circle **Y** (Yes) if it was a concern for you or circle **N** (No) if it was not a concern for you.

- |                                       | No | Yes |
|---------------------------------------|----|-----|
| a. Some shots are given too early     | N  | Y   |
| b. Too many shots are given at a time | N  | Y   |

- c. I do not feel some of the diseases will affect my child N Y
- d. Shots may weaken my child's immune system N Y
- e. Some of the shots do more harm than good N Y
- f. Getting some of the childhood diseases is natural N Y
- g. I have religious beliefs or concerns about SOME shots N Y
- h. I have religious beliefs or concerns about ALL shots N Y
- i. Other → Please tell us: N Y

**74. The flu season in Oregon usually runs September thru March of each year. Has your two-year-old ever had a flu vaccination or shot anytime during:**

- |                                                                               | No | Yes | Don't Know |
|-------------------------------------------------------------------------------|----|-----|------------|
| a. This year's flu season (September thru March of <i>this</i> calendar year) | N  | Y   | DK         |
| b. Last year's flu season (September thru March of <i>last</i> calendar year) | N  | Y   | DK         |

**75. Has a health care provider ever given an immunization or baby shot to your two-year-old during a sick or urgent care visit?**

- No
- Yes

*The next questions are about your two-year-old's dental care.*

**76. Has your two-year-old ever been to a dentist or dental clinic?**

- No
- Yes → Go to Question 78

**77. Here is a list of problems some people can have getting dental care for their children.** For each item, circle Y (Yes) if it was a problem for you or circle N (No) if it was not a problem or did not apply to you.

- |                                                                            | No | Yes |
|----------------------------------------------------------------------------|----|-----|
| a. I didn't have enough money to pay for the visit                         | N  | Y   |
| b. I didn't have insurance to pay for the visit                            | N  | Y   |
| c. I couldn't locate a dentist who would see my child                      | N  | Y   |
| d. I couldn't get an appointment with a dentist                            | N  | Y   |
| e. A health care or dental care provider told me my child was too young to |    |     |

- see the dentist N Y
- f. I didn't think my child needed to go N Y
- g. I had no one to take care of my other children N Y
- h. I had too many other things going on N Y
- i. Other → Please tell us: N Y

**78. Does your two-year-old receive fluoride drops or tablets daily?**

- No
- Yes

*The next questions are about your two-year-old's medical history.*

**79. Please circle Y (Yes) or N (No) for each of the following.**

**Does your two-year-old have . . . ?**

**a. A diagnosis of a chronic condition such as:**

- |                             | No | Yes |
|-----------------------------|----|-----|
| (1) Asthma                  | N  | Y   |
| (2) Autism                  | N  | Y   |
| (3) Cleft palate            | N  | Y   |
| (4) Down syndrome           | N  | Y   |
| (5) Cerebral palsy          | N  | Y   |
| (6) Other chronic condition | N  | Y   |

Please tell us:

**b. An ongoing need (lasting six months or more) for:**

- |                                                 |   |   |
|-------------------------------------------------|---|---|
| (1) Specialty health care                       | N | Y |
| (2) Behavioral health or mental health services | N | Y |
| (3) Physical therapy                            | N | Y |
| (4) Occupational therapy                        | N | Y |
| (5) Speech services                             | N | Y |

**c. An ongoing need (lasting six months or more) for:**

- |                               |   |   |
|-------------------------------|---|---|
| (1) Medication                | N | Y |
| (2) Home health services      | N | Y |
| (3) Special diet              | N | Y |
| (4) Use of assistive devices  | N | Y |
| (5) Durable medical equipment | N | Y |

**80. Please circle Y (Yes) or N (No) for each of the following.**

**Does your two-year-old.....?**

- |                                                                                           | No | Yes |
|-------------------------------------------------------------------------------------------|----|-----|
| b. Need more time at doctor's visits than usual for children his/her age                  | N  | Y   |
| c. Need more frequent office visits than usual for children his/her age                   | N  | Y   |
| d. Need or use more medical or mental health services than usual for children his/her age | N  | Y   |
| e. Currently need or use medicine (other                                                  |    |     |

- than vitamins) prescribed by a doctor N Y
- f. Seem limited or prevented in any way in his or her ability to do the things most two-year-olds can do N Y
- g. Experience any kind of emotional, developmental or behavioral problem for which he/she needs treatment or counseling N Y

**81. Early Intervention Services is a State program that offers free services to children age 3 and under who have developmental problems or delays. Has your two-year-old ever....?**

- |                                                                    | No | Yes |
|--------------------------------------------------------------------|----|-----|
| a. Been referred for Early Intervention Service                    | N  | Y   |
| b. Been screened or tested for Early Intervention services         | N  | Y   |
| c. Been found eligible (qualified) for Early Intervention services | N  | Y   |
| d. Received Early Intervention services                            | N  | Y   |

**If your two-year-old has never been referred for Early Intervention, please go to Question 83.**

**82. Below are reasons why children who were referred for Early Intervention may not receive services. For each item, circle Y (Yes) if it was a reason for your two-year-old or circle N (No) if it was not.**

- |                                                                                      | No | Yes |
|--------------------------------------------------------------------------------------|----|-----|
| a. I don't think my child needs Early Intervention services                          | N  | Y   |
| b. My child is getting private services instead                                      | N  | Y   |
| c. I don't know how to get my child tested                                           | N  | Y   |
| d. The testing process is too confusing and complicated                              | N  | Y   |
| e. My child was tested but not found eligible                                        | N  | Y   |
| f. My child was tested and is eligible. We have been waiting ___ months for services | N  | Y   |
| g. There are no openings right now                                                   | N  | Y   |
| h. I can't get time off to take my child                                             | N  | Y   |
| i. I don't have childcare for my other kids and can't take them with us              | N  | Y   |
| j. I don't have transportation                                                       | N  | Y   |
| k. We moved                                                                          | N  | Y   |
| l. Other → Please tell us:                                                           | N  | Y   |

***The next questions are about your two-year-old's current activities.***

**83. Do you have regular childcare arrangements for your two-year-old now?**

- No → Go to Page 14, Question 86

- Yes

**84. What are your childcare arrangements? Check all that apply.**

- Childcare in non-relative's home
- Childcare center
- Paid care in your home
- Other older children
- Child's grandparent(s)
- Other relative(s)
- Baby-sitter/friend/neighbor
- Other → Please tell us:

**85. What is the average number of hours per week that your two-year-old stays in childcare?**

- Less than 10 hours per week
- 10 to 19 hours per week
- 20 to 29 hours per week
- 30 to 39 hours per week
- 40 hours or more per week

**86. In a typical day, how much time does your two-year-old spend watching TV or videos? Check one answer.**

- None
- Less than 2 hours
- 2 hours or more

**87. Are you concerned about the amount of TV your two-year-old watches?**

- No
- Yes

**88. In a typical week, how often do you, or someone else in your household, read a book or story to your two-year-old? Check one answer.**

- Every day
- At least three times a week
- Once a week
- Less than once a week
- Never

**89. How many times in the past week have you or any family member taken your two-year-old on any kind of outing, such as to a park, playground, library or other children's program or activity?**

- None
- 1 to 3 times
- 4 to 5 times
- 6 or more times

**90. About how many hours a day, on average, is your two-year-old in the same room with someone who is smoking?**

\_\_\_\_\_ Hours

- Less than 1 hour a day
- My two-year-old is never in the same room with someone who is smoking

**91. Is there a TV in your two-year-old's bedroom?**

- No
- Yes

**92. What is today's date?**

\_\_\_\_\_ 20\_\_\_\_\_  
Month Day Year

*Thank you for taking the time to answer these questions. Your answers are important and could help us learn about ways to improve the health of children in the future.*

**Please use this space for any additional comments you would like to make about the health of mothers and their children in Oregon.**

## Appendix C

### Calculation of un-weighted response rate for PRAMS-2:

Number of PRAMS-2 respondents/Total number of women who responded to PRAMS  
= 865/1,968 = 44%

### Calculation of weighted response rate according to the formula identified by the CDC:

$\sum$  PRAMS weights (of PRAMS-2 responders) /  $\sum$  PRAMS weights (total)  
= 22535.71/44131 = 51.1%

### Calculation of overall response rate for both PRAMS & PRAMS-2 combined:

Response rate for PRAMS \* Response rate for PRAMS-2  
= 0.748 \* 0.511 = 38.2%

### Verification of response rate calculations with TOTS

Since 1994, the state of Oklahoma's Maternal & Child Health department has a survey called TOTS (The Oklahoma Toddler Survey), which is a 2-year longitudinal follow-up of Oklahoma PRAMS respondents. This is similar to what the state of Oregon started doing starting in 2006. Thus, calculation of weighted response rates for PRAMS-2 and overall response rates for PRAMS & PRAMS-2 were verified with the Oklahoma state department of Maternal & Child Health. Below is the actual correspondence from Robert W. Feyerharm (Oklahoma State department of Maternal and Child Health) regarding calculation of calculating a weighted response rate for PRAMS-2.

---

So the steps should be:

1. Sum the weights of only those women who responded to PRAMS2.
2. Sum the weights of all women (responders & nonresponders) in the PRAMS1 sample.
3. Divide 1 by 2.

The 69% weighted response rate tells you the adjusted % of women who responded to PRAMS2, given that they had already responded to PRAMS1. However, since PRAMS2 was sampled from PRAMS1 responders, it makes more sense to report the weighted response rate of 51.1%, which is the % of women in the original PRAMS1 sampling frame who completed both PRAMS1 and PRAMS2.

One of the issues with longitudinal studies is that the overall response rate may become progressively smaller with each successive survey. We have response rates of about 70% in Oklahoma for PRAMS and TOTS, so that only around 50% (.70 x .70 = .49) of the original mothers in the PRAMS sample respond to both PRAMS and TOTS.

Robert

---

### Verification of response rate calculation with Alaska CUBS

Alaska CUBS (Childhood Understanding Behaviors Survey) is a follow-up survey to PRAMS sent to PRAMS mothers who now have a 3-year old child. Methodology for response rate calculations was found to be similar to the CDC and Oklahoma state

methodologies. Below is the actual correspondence from Margaret Blabey, MPH (CUBS Program Coordinator and Infant Mortality Epidemiologist).

---

We are just now getting our 2008 CUBS data weighted so I do not have a weighted response rate for that yet. Our unweighted response rate for last year was 49%.

For our 2006 data (our first year of CUBS), our weighted response rate was 53.7% (unweighted 50.7%). The denominator for this was all women who were eligible for CUBS (so, they returned a contact card with PRAMS in 2004 AND were not dropped due to any reason other than mom died or mom incapable of doing the survey). To calculate the weighted response rate, I weighted the data by WTONE (sample weight) and ran a frequency of INQX, within the subpopulation of eligible moms.

Margaret

---

### **Verification of response rate calculation with MEPS (Medical Expenditure Panel Survey)**

The Medical Expenditure Panel Survey (MEPS) is a set of large-scale surveys of families and individuals, their medical providers, and employers across the United States. MEPS is the most complete source of data on the cost and use of health care and health insurance coverage. The panel design of the survey, which features several rounds of interviewing covering two full calendar years, makes it possible to examine changes in different health care related variables over time.

MEPS uses a multiplicative function of the wave or round-specific response rates to calculate the overall response rate. Detailed methodology on how overall response rates for MEPS are calculated is given in the document titled 'Medical Expenditure Panel Survey Household Component'<sup>47</sup>.

## References

1. Adams, E.K., Miller, V.P., Ernst, C., Nishimura, B.K., Melvin, C, Merritt, R. (2002). Neonatal health care costs related to smoking during pregnancy. *Health Economics*, 11, 193-206.
2. Ahluwalia, I.B., Mack, K.A. & Mokdad, A. (2004). Mental and physical distress and high-risk behaviors among reproductive-age women. *American College of Obstetricians and Gynecologists*, 104(3), 477-483.
3. Allen, A.M., Prince, C.B., Dietz, P.M. (2009). Postpartum depressive symptoms and smoking relapse. *American Journal of Preventive Medicine*, 36(1), 9-12.
4. American Cancer Society, Inc. (2008). *Smoking costs US \$157 billion each year*. Retrieved November 10, 2008 from [http://www.cancer.org/docroot/NWS/content/NWS\\_1\\_1x\\_Smoking\\_Costs\\_US\\_157\\_Billion\\_Each\\_Year.asp](http://www.cancer.org/docroot/NWS/content/NWS_1_1x_Smoking_Costs_US_157_Billion_Each_Year.asp)
5. Armour, B.S., Pitts, M.M., Lee, C.W. (2008). Cigarette smoking and food insecurity among low-income families in the United States, 2001. *American Journal of Health Promotion*, 22(6), 386-392. [Abstract retrieved from PubMed – 2/22/2009].
6. Asomaning, K., Miller, D.P., Liu, G., Wain, J.C., Lynch, T.J., Su, L., et al. (2008). Second hand smoke, age of exposure and lung cancer risk [Abstract]. *Lung Cancer*, 61(1), 13-20. Retrieved from PubMed database.
7. Bottorff, J.L., Johnson, J.L., Irwin, L.G. & Ratner, P.A. (2000). Narratives of smoking relapse: The stories of postpartum women. *Research in Nursing & Health*, 23, 126-134.
8. Campaign for Tobacco-Free Kids. (2008). *The Toll of Tobacco*. Retrieved on November 10, 2008 from <http://www.tobaccofreekids.org/reports/settlements/toll.php?StateID=OR>
9. Carmichael, S.L., Ahluwalia, I.B., & the PRAMS working group. (2000). Correlates of postpartum smoking relapse – Results from PRAMS. *American Journal of Preventive Medicine*, 19(3), 193-196.
10. Centers for Disease Control and Prevention (CDC). (2007). *Pregnancy Risk Assessment Monitoring System (PRAMS): Methodology*. Retrieved on November 10, 2008 from <http://www.cdc.gov/PRAMS/methodology.htm>



11. Chaloupka, F.J. (2008). Smoking, food insecurity, and tobacco control. *Archives of Pediatric Adolescent Medicine*, 162(11).
12. Cigarette Smoking During the Last 3 Months of Pregnancy Among Women Who Gave Birth to Live Infants--Maine, 1988-1997.. (1999, May 28). *MMWR: Morbidity & Mortality Weekly Report*, Retrieved February 22, 2009, from Health Source: Nursing/Academic Edition database.
13. Clark, E. & Rosenberg, K.D. (2003). Risk factors for smoking cessation relapse after pregnancy. *Proceedings from the 9<sup>th</sup> Annual Maternal and Child Health Epidemiology Workshop*. Tempe, AZ.
14. Colman, G.J. & Joyce, T. (2003). Trends in Smoking Before, During and After Pregnancy in Ten States. *American Journal of Preventive Medicine*, 24(1), 29-35.
15. Davis, C.S. (2002). *Statistical methods for the analysis of repeated measurements*. Springer-Verlag: New York.
16. Der, G. & Everitt, B.S. (2006). *Statistical Analysis of Medical Data Using SAS*. Chapman & Hall: Boca Raton.
17. Ebert, L.M., Fahy, K. (1993). Why do women continue to smoke in pregnancy? *The Journal of the Oklahoma State Medical Association*, 86(5), 231-236.
18. Gaffney, K.F. & Henry, L.L. (2007). Identifying risk factors for postpartum tobacco use. *Journal of Nursing Scholarship*, 39(2), 126-132.
19. Gaffney, K.F., Beckwitt, A.E. & Friesen, M.A. (2008). Mothers' reflections about infant irritability and postpartum tobacco use [Abstract]. *BIRTH*, 35(1). Retrieved from PubMed database.
20. Hensley, A.S.M., Lappin, R.E., Peterson, L., Johnson, C.C. (2007). Pregnancy associated smoking behavior and six year postpartum recall. *Women Birth*, 20(4), 161-168.
21. Kahn, R.S., Certain, L. & Whitaker, R.C. (2002). A reexamination of smoking before, during, and after pregnancy. *American Journal of Public Health*, 92(11), 1801-1808.
22. Kost, K., Landry, D.J., Darroch, J.E. (1998). Predicting maternal behaviors during pregnancy: does intention status matter? *Family Planning Perspectives*, 30(2), 79-88.

23. Lelong, N., Kaminski, M., Saurel-Cubizolles, M. & Bouvier-Colle, M. (2001). Postpartum return to smoking among usual smokers who quit during pregnancy. *European Journal of Public Health*, 11(3), 334-339.
24. Levine, M.D. & Marcus, M.D. (2004). Do changes in mood and concerns about weight relate to smoking relapse in the postpartum period? *Archives of Women's Mental Health*, 7, 155-156.
25. Levine, M.D., Marcus, M.D. & Kalarchian, M.A. (2006). Weight concerns affect motivation to remain abstinent from smoking postpartum. *Annals of Behavioral Medicine*, 32(2), 147-153.
26. Liu, J., Rosenberg, K.D. & Sandoval, A.P. (2006). Breastfeeding duration and perinatal cigarette smoking in a population-based cohort. *American Journal of Public Health*, 96(2), 309-314.
27. McBride, C.M., Curry, S.J., Lando, H.A., Pirie, P.L., Grothuas, L.C. & Nelson, J.C. (1999). Prevention of relapse in women who quit smoking during pregnancy. *American Journal of Public Health*, 89(5), 706-711.
28. Mullen, P.D., Richardson, M.A., Quinn, V.P. & Ershoff, D.H. (1997). Postpartum return to smoking: Who is at risk and when [Abstract]. *American Journal of Health Promotion*, 11(5). Retrieved from PubMed database.
29. O'Campo, P., Faden, R.R., Brown, H. & Gielen, A.C. (1992). The impact of pregnancy on women's prenatal and postpartum smoking behavior [Abstract]. *American Journal of Preventive Medicine*, 8(1). Retrieved from PubMed database.
30. Prochaska, J.O. & Velicer, W.F. (2004). Integrating population smoking cessation policies and programs. *Public Health Reports*, 119, 244-252.
31. Ramsey, A.M., Blose, D., Lorenz, D., Thomas, W., DePersio, S.R., Bruce, F.C. (1993). Cigarette smoking among women in Oklahoma: before, during, and after pregnancy. *Journal of Oklahoma State Medical Association*, 86(5), 231-236.
32. Reitzel, L.R., Vidrine, J.I., Yisheng, L., Mullen, P.D., Velasquez, M.M., Cinciripini, P.M. et al. (2007). The influence of subjective social status on vulnerability to postpartum smoking among young pregnant women. *American Journal of Public Health*, 97(8), 1476-1482.

33. Repko, M., van den Brink, W., Huyser, J. (2007). Can depressed patients stop smoking? A review. *Dutch Journal of Psychiatry*, 49(10), 753-762. [Abstract accessed from PubMed on April 9, 2009].
34. Ripley-Moffitt, C.E. (2008). *Newly-defined factors may prevent postpartum smoking relapse*. Retrieved November 5, 2008 from <http://www.unchealthcare.org/site/newsroom/news/2008/August/pregnancysmoking/>
35. Severson, H.H., Andrews, J.A., Lichtenstein, E., Wall, M. & Zoref, L. (1995). Predictors of smoking during and after pregnancy: A survey of mothers of newborns. *Preventive Medicine*, 24, 23-28.
36. Rosenberg, K.D., Gelow, J.M. & Sandoval, A.P. (2003). Pregnancy intendedness and the use of periconceptional folic acid. *Pediatrics*, 111(5), 1142-1145. Retrieved November 8, 2008 from <http://pediatrics.aappublications.org/cgi/content/full/111/5/S1/1142>
37. Solomon, L.J., Higgins, S.T., Heil, S.H., Badger, G.J., Thomas, C.S. & Bernstein, I.M. (2007). Predictors of postpartum relapse to smoking [Abstract]. *Drug Alcohol Dependency*, 90, 2-3. Retrieved from PubMed database.
38. Tager, I.B. (2008). The effects of second-hand and direct exposure to tobacco smoke on asthma and lung function in adolescence [Abstract]. *Pediatric Respiratory Reviews*, 9(1), 29-37. Retrieved from PubMed database.
39. Tong, V.T., England, L.J., Dietz, P.M., Asare, A.L. (2008). Smoking patterns and use of cessation interventions during pregnancy. *American Journal of Preventive Medicine*, 35(4), 327-333.
40. U.S. Census Bureau: State and County QuickFacts (last updated Feb 20, 2009). Retrieved April 14, 2008, from <http://quickfacts.census.gov/qfd/states/41000.html>
41. Whitaker, R.C., Orzol, S.M. & Kahn, R.S. (2007). The co-occurrence of smoking and a major depressive episode among mothers 15 months after delivery. *Preventive Medicine*, 45, 476-480.
42. Yates, J.T. (2008). Maternal depression and food insecurity during pregnancy among Oregon women. Thesis presented to the department of Public Health and Preventive Medicine of Oregon Health and Science University (OHSU) in partial fulfillment of the requirements of the degree of Master of Public Health.

43. Yu, S.M., Park, C.H. & Schwalberg, R.H. (2002). Factors Associated with Smoking Cessation Among U.S. Pregnant Women. *Maternal and Child Health Journal*, 6(2), 89-97.
44. Information about Oregon's safety net programs. (2008). Retrieved on March 30, 2009 from [http://oregonstate.edu/instruct/dce/chi/information\\_or\\_safety.html](http://oregonstate.edu/instruct/dce/chi/information_or_safety.html)
45. Hughes JR, Stead LF, Lancaster T. Antidepressants for smoking cessation. *Cochrane Database of Systematic Reviews* 2007, Issue 1. Art. No.: CD000031. DOI: 10.1002/14651858.CD000031.pub3.
46. Hosmer, D.W., Lemeshow, S. (2000). *Applied Logistic Regression: Second Edition*. John Wiley & Sons, Canada.
47. Medical Expenditure Panel Survey Household Component: Public Use File 1, retrieved on May 5, 2009 from [http://www.meps.ahrq.gov/mepsweb/data\\_stats/download\\_data/pufs/h01/h01doc.pdf#xml=http://meps.ahrq.gov/cgi-bin/taxis/webinator/search/pdfhi.txt?query=calculating+response+rate&pr=MEPS\\_FULLSITE&prox=page&rorder=500&rprox=500&rdfreq=500&rwfreq=500&rlead=500&sufs=0&order=r&cq=&id=49feef3310](http://www.meps.ahrq.gov/mepsweb/data_stats/download_data/pufs/h01/h01doc.pdf#xml=http://meps.ahrq.gov/cgi-bin/taxis/webinator/search/pdfhi.txt?query=calculating+response+rate&pr=MEPS_FULLSITE&prox=page&rorder=500&rprox=500&rdfreq=500&rwfreq=500&rlead=500&sufs=0&order=r&cq=&id=49feef3310).